

FIRST INTEGRALS OF LINEAR DIFFERENTIAL SYSTEMS

V.N.Gorbuzov*, A.F. Pranevich**

**Department of Mathematics and Computer Science, Yanka Kupala Grodno State University,
Ozeshko 22, Grodno, 230023, Belarus*

E-mail: gorbuzov@grsu.by

***Department of Economics and Management, Yanka Kupala Grodno State University,
Ozeshko 22, Grodno, 230023, Belarus*

E-mail: pronevich@tut.by

Abstract

We investigate the problem of the existence of first integrals for multidimensional and ordinary linear differential systems with constant coefficients. The spectral method of the first integrals basis construction for these systems of linear differential equations is developed.

Key words: linear system of equations in total differentials, ordinary linear differential system, partial integral, first integral.

2000 Mathematics Subject Classification: 34A30

Contents

1. $\mathbb{R}$-differentiable integrals of $\mathbb{R}$-linear systems in total differentials	2
1.1. $\mathbb{R}$ -linear homogeneous systems in total differentials ¹	2
1.1.1. $\mathbb{R}$ -linear partial integral	2
1.1.2. Autonomous $\mathbb{R}$ -differentiable first integrals	3
1.1.3. Nonautonomous $\mathbb{R}$ -differentiable first integrals	7
1.2. $\mathbb{R}$ -linear nonhomogeneous systems in total differentials	8
2. First integrals of linear real systems in total differentials	9
2.1. Linear real homogeneous systems in total differentials ²	9
2.1.1. Partial integrals	9
2.1.2. Autonomous first integrals	10
2.1.3. Nonautonomous first integrals	20
2.2. Linear real nonhomogeneous systems in total differentials	22
3. First integrals of linear systems of ordinary differential equations³	26
3.1. Linear homogeneous systems of ordinary differential equations	26
3.1.1. Autonomous first integrals	26
3.1.2. Nonautonomous first integrals	30
3.2. Linear nonhomogeneous systems of ordinary differential equations	32
References	34

¹This Subsection has been published in *Dokl. Akad. Nauk Belarusi* 48 (2004), No. 1, 49-52.

²The definitive version of this Subsection has been published in *Differential equations and control processes*, 2001, No.3, 17-45 (<http://www.neva.ru/journal>); Deponent VINITI of *Differentsial Uravneniya (Differential Equations)*, 02.10.2002, No. 1667-B2002; *Mathematical research*, Vol. 10 (2003), 143-152.

³Section 3 has been published in *Vestnik of the Yanka Kupala Grodno State Univ.*, 2002, Ser. 2, No. 2(11), 23-29; 2003, Ser. 2, No. 2(22), 50-60; 2008, Ser. 2, No. 2(68), 5-10.

1. $\mathbb{R}$ -differentiable integrals of $\mathbb{R}$ -linear systems in total differentials

1.1. $\mathbb{R}$ -linear homogeneous systems in total differentials

We consider the system of equations in total differentials

$$dw = X_1(w)dz + X_2(w)d\bar{z}, \quad (1.1)$$

where $w = \text{colon}(w_1, \dots, w_n) \in \mathbb{C}^n$, $z = \text{colon}(z_1, \dots, z_m) \in \mathbb{C}^m$; $dw = \text{colon}(dw_1, \dots, dw_n)$, $dz = \text{colon}(dz_1, \dots, dz_m)$, and $d\bar{z} = \text{colon}(d\bar{z}_1, \dots, d\bar{z}_m)$ are vector columns; $\bar{z}_j$ is complex conjugate of z_j ; the entries of the matrices $X_1(w) = \|X_{\tau j}(w)\|$ and $X_2(w) = \|X_{\tau, m+j}(w)\|$, $\tau = 1, \dots, n$, $j = 1, \dots, m$ are the $\mathbb{R}$ -linear functions [1, p. 21]

$$X_{\tau k}: w \rightarrow \sum_{\xi=1}^n (a_{\tau k \xi} w_{\xi} + a_{\tau k, n+\xi} \bar{w}_{\xi}) \quad \text{for all } w \in \mathbb{C}^n, \quad k = 1, \dots, 2m, \quad \tau = 1, \dots, n,$$

with constant coefficients $a_{\tau k \varrho} \in \mathbb{C}$, $\varrho = 1, \dots, 2n$, $k = 1, \dots, 2m$, $\tau = 1, \dots, n$. We assume that the linear differential operators

$$\mathfrak{x}_j(w) = \sum_{\xi=1}^n (X_{\xi j}(w) \partial_{w_{\xi}} + \bar{X}_{\xi, m+j}(w) \partial_{\bar{w}_{\xi}}) \quad \text{for all } w \in \mathbb{C}^n, \quad j = 1, \dots, m, \quad (1.2)$$

and

$$\mathfrak{x}_{m+j}(w) = \sum_{\xi=1}^n (X_{\xi, m+j}(w) \partial_{w_{\xi}} + \bar{X}_{\xi j}(w) \partial_{\bar{w}_{\xi}}) \quad \text{for all } w \in \mathbb{C}^n, \quad j = 1, \dots, m, \quad (1.3)$$

induced by this system are related by the Frobenius conditions [2; 3]. These conditions are represented via Poisson brackets as the system of identities

$$[\mathfrak{x}_k(w), \mathfrak{x}_l(w)] = \mathfrak{D} \quad \text{for all } w \in \mathbb{C}^n, \quad k = 1, \dots, 2m, \quad l = 1, \dots, 2m, \quad (1.4)$$

i.e., system (1.1) is completely solvable [4; 5, pp. 15 – 25].

A general integral of the completely solvable system in total differentials (1.1) is n functionally independent $\mathbb{R}$ -differentiable first integrals of (1.1). The completely solvable differential system (1.1) has also $n - m$ autonomous $\mathbb{R}$ -differentiable first integrals (see [6]).

In this paper we study Darboux's problem of finding first integrals in case that partial integrals are known [7]. Using method of partial integrals [5; 8; 9], we obtain the spectral method for building first integrals of linear differential systems [5, pp. 239 – 272; 10 – 17].

The problems related to those studied in the present paper were intensively investigated by many people. For example, see for systems of ordinary differential equations [7; 12; 16 – 54], for partial differential systems [5; 8; 10; 15; 55 – 59], for systems of equations in total differentials [2 – 6; 8; 9; 11; 13; 14; 16; 32; 60 – 63], and this list is very far from being complete.

1.1.1. $\mathbb{R}$ -linear partial integral. The $\mathbb{R}$ -linear function

$$p: w \rightarrow \sum_{\xi=1}^n (b_{\xi} w_{\xi} + b_{n+\xi} \bar{w}_{\xi}) \quad \text{for all } w \in \mathbb{C}^n \quad (b_{\varrho} \in \mathbb{C}, \quad \varrho = 1, \dots, 2n)$$

is a *partial integral* of the system in total differentials (1.1) iff

$$\mathfrak{x}_k p(w) = p(w) \lambda^k \quad \text{for all } w \in \mathbb{C}^n, \quad \lambda^k \in \mathbb{C}, \quad k = 1, \dots, 2m.$$

This system of identities is equivalent to the linear homogeneous system

$$(A_k - \lambda^k E) b = 0, \quad k = 1, \dots, 2m, \quad (1.5)$$

where $A_j = \|a_{1j} \dots a_{nj} \bar{a}_{1, m+j} \dots \bar{a}_{n, m+j}\|$ and $A_{m+j} = \|a_{1, m+j} \dots a_{n, m+j} \bar{a}_{1j} \dots \bar{a}_{nj}\|$ are the

$2n \times 2n$ -matrices with $a_{\tau k} = \text{colon}(a_{\tau k 1}, \dots, a_{\tau k, 2n})$, $\bar{a}_{\tau k} = \text{colon}(\bar{a}_{\tau k, n+1}, \dots, \bar{a}_{\tau k, 2n}, \bar{a}_{\tau k 1}, \dots, \bar{a}_{\tau k n})$, $\tau = 1, \dots, n$, $k = 1, \dots, 2m$, $j = 1, \dots, m$, E is the $2n \times 2n$ identity matrix, and $b = \text{colon}(b_1, \dots, b_{2n})$ is a vector column.

The Frobenius conditions (1.4) for system (1.1) are equivalent [60, p. 73]

$$A_k A_l = A_l A_k, \quad k = 1, \dots, 2m, \quad l = 1, \dots, 2m.$$

Then there exists a relation [64, pp. 193 – 194; 65] between eigenvectors and eigenvalues of the matrices A_k , $k = 1, \dots, 2m$.

Lemma 1.1. *Let $\nu \in \mathbb{C}^{2n}$ be a common eigenvector of the matrices A_k , $k = 1, \dots, 2m$. Then the $\mathbb{R}$ -linear function*

$$p: w \rightarrow \nu \gamma \quad \text{for all } w \in \mathbb{C}^n,$$

where $\gamma = \text{colon}(w_1, \dots, w_n, \bar{w}_1, \dots, \bar{w}_n)$, is a partial integral of the system (1.1).

Proof. If ν is a common eigenvector of the matrices A_k , $k = 1, \dots, 2m$, then ν is a solution to system (1.5), where λ^k is an eigenvalue of the matrix A_k corresponding to the eigenvector ν . We obtain $\mathfrak{x}_k(\nu \gamma) = \lambda^k \nu \gamma$ for all $w \in \mathbb{C}^n$, $k = 1, \dots, 2m$. Therefore the $\mathbb{R}$ -linear function p is a partial integral of the system in total differential (1.1). ■

1.1.2. Autonomous $\mathbb{R}$ -differentiable first integrals

Theorem 1.1. *Let ν^θ , $\theta = 1, \dots, 2m+1$ be common eigenvectors of the matrices A_k , $k = 1, \dots, 2m$. Then the system (1.1) has the $\mathbb{R}$ -differentiable autonomous first integral*

$$F: w \rightarrow \prod_{\theta=1}^{2m+1} (\nu^\theta \gamma)^{h_\theta} \quad \text{for all } w \in \Omega, \quad \Omega \subset D(F), \quad (1.6)$$

where $h_1, \dots, h_{2m+1}$ is a nontrivial solution to the system $\sum_{\theta=1}^{2m+1} \lambda_\theta^k h_\theta = 0$, $k = 1, \dots, 2m$, and λ_θ^k are the eigenvalues of the matrices A_k , $k = 1, \dots, 2m$, corresponding to the common eigenvectors ν^θ , $\theta = 1, \dots, 2m+1$.

Proof. Suppose ν^θ are common eigenvectors of the matrices A_k corresponding to the eigenvalues λ_θ^k , $k = 1, \dots, 2m$, $\theta = 1, \dots, 2m+1$, respectively. By lemma 1.1, it follows that the $\mathbb{R}$ -linear functions $w \rightarrow \nu^\theta \gamma$ for all $w \in \mathbb{C}^n$, $\theta = 1, \dots, 2m+1$ are partial integrals of the system of equations in total differentials (1.1). Hence,

$$\mathfrak{x}_k \nu^\theta \gamma = \lambda_\theta^k \nu^\theta \gamma \quad \text{for all } w \in \mathbb{C}^n, \quad k = 1, \dots, 2m, \quad \theta = 1, \dots, 2m+1. \quad (1.7)$$

We form the function

$$F: w \rightarrow \prod_{\theta=1}^{2m+1} (\nu^\theta \gamma)^{h_\theta} \quad \text{for all } w \in \Omega, \quad \Omega \subset \mathbb{C}^n,$$

where Ω is a domain (open arcwise connected set) in $\mathbb{C}^n$ and h_θ , $\theta = 1, \dots, 2m+1$ are complex numbers with $\sum_{\theta=1}^{2m+1} |h_\theta| \neq 0$. The Lie derivative of F by virtue of (1.1) is equal to

$$\mathfrak{x}_k F(w) = \prod_{\theta=1}^{2m+1} (\nu^\theta \gamma)^{h_\theta-1} \sum_{\theta=1}^{2m+1} h_\theta \prod_{l=1, l \neq \theta}^{2m+1} (\nu^l \gamma) \mathfrak{x}_k \nu^\theta \gamma \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m.$$

Using (1.7), we get

$$\mathfrak{x}_k F(w) = \sum_{\theta=1}^{2m+1} \lambda_\theta^k h_\theta F(w) \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m.$$

If $\sum_{\theta=1}^{2m+1} \lambda_{\theta}^k h_{\theta} = 0$, $k = 1, \dots, 2m$, then the function (1.6) is an autonomous $\mathbb{R}$ -differentiable first integral of the system in total differentials (1.1). ■

Corollary 1.1. *Let ν^{θ} be common eigenvectors of the matrices A_k corresponding to the eigenvalues λ_{θ}^k , $k = 1, \dots, 2m$, $\theta = 1, \dots, 2m+1$, respectively. Then the system of equations in total differentials (1.1) has the $\mathbb{R}$ -differentiable autonomous first integral*

$$F_{12\dots 2m(2m+1)} : w \rightarrow \prod_{\theta=1}^{2m} (\nu^{\theta} \gamma)^{-\delta_{\theta}} (\nu^{2m+1} \gamma)^{\delta} \quad \text{for all } w \in \Omega, \quad \Omega \subset D(F_{12\dots 2m(2m+1)}),$$

where the determinants δ_{θ} , $\theta = 1, \dots, 2m$ are obtained by replacing the θ -th column of the determinant $\delta = |\lambda_{\theta}^k|$ by $\text{colon}(\lambda_{2m+1}^1, \dots, \lambda_{2m+1}^{2m})$, respectively.

For example, the $\mathbb{R}$ -linear autonomous system of equations in total differentials

$$\begin{aligned} dw_1 &= (2w_1 - i(w_2 + \bar{w}_1) + (1-i)\bar{w}_2)dz + (w_1 + (2-i)(w_2 + \bar{w}_1) + (1-i)\bar{w}_2)d\bar{z}, \\ dw_2 &= -((2+i)(w_1 + \bar{w}_2) + iw_2)dz - (i(w_1 + \bar{w}_2) + (i-1)w_2)d\bar{z} \end{aligned} \quad (1.8)$$

has the commuting matrices

$$A_1 = \begin{vmatrix} 2 & -2-i & 2+i & 0 \\ -i & -i & 1+i & i \\ -i & 0 & 1 & i \\ 1-i & -2-i & 2+i & 1+i \end{vmatrix} \quad \text{and} \quad A_2 = \begin{vmatrix} 1 & -i & i & 0 \\ 2-i & 1-i & 1+i & -2+i \\ 2-i & 0 & 2 & -2+i \\ 1-i & -i & i & i \end{vmatrix}.$$

Therefore the system of equations in total differentials (1.8) is completely solvable.

The matrices A_1 and A_2 have the eigenvalues $\lambda_1^1 = 1+i$, $\lambda_2^1 = -i$, $\lambda_3^1 = 1$, $\lambda_4^1 = 2$, and $\lambda_1^2 = i$, $\lambda_2^2 = 1-i$, $\lambda_3^2 = 2$, $\lambda_4^2 = 1$ corresponding to the eigenvectors $\nu^1 = (0, 1, 1, 1)$, $\nu^2 = (1, 1, 0, 1)$, $\nu^3 = (0, 1, 1, 0)$, $\nu^4 = (1, 0, 0, 1)$, respectively.

The solution to the linear homogeneous system

$$\begin{cases} (1+i)h_1 - ih_2 + h_3 = 0, \\ ih_1 + (1-i)h_2 + 2h_3 = 0 \end{cases} \iff \begin{cases} h_1 = -(1+i)h_3, \\ h_2 = -(2+i)h_3 \end{cases}$$

is $h_1 = 1+i$, $h_2 = 2+i$, $h_3 = -1$.

The $\mathbb{R}$ -differentiable function (by Theorem 1.1)

$$F : w \rightarrow \frac{(w_2 + \bar{w}_1 + \bar{w}_2)^{1+i}(w_1 + w_2 + \bar{w}_2)^{2+i}}{w_2 + \bar{w}_1} \quad \text{for all } w \in \Omega, \quad (1.9)$$

where a domain $\Omega \subset \{w : w_2 + \bar{w}_1 \neq 0\}$, is an autonomous first integral of the system (1.8).

The $\mathbb{R}$ -differentiable first integral (1.9) is an autonomous general integral of the completely solvable system of equations in total differentials (1.8).

From the entire set of ordinary differential systems induced by the completely solvable system of equations in total differentials (1.1), we extract system

$$dw_{\tau} = X_{\tau\zeta}(w) dz_{\zeta} + X_{\tau, m+\zeta}(w) d\bar{z}_{\zeta}, \quad \tau = 1, \dots, n, \quad \zeta \in \{1, \dots, m\} \quad (1.1.\zeta)$$

such that the matrix A_{ζ} has the smallest number of elementary divisors [64, p. 147].

Definition 1.1. *Let ν^{0l} be an eigenvector of the matrix A_{ζ} corresponding to the eigenvalue λ_l^{ζ} with elementary divisor of multiplicity s_l . A non-zero vector $\nu^{\eta l} \in \mathbb{C}^{2n}$ is called a **generalized eigenvector of order η** for λ_l^{ζ} if and only if*

$$(A_{\zeta} - \lambda_l^{\zeta} E) \nu^{\eta} = \eta \cdot \nu^{\eta-1}, \quad \eta = 1, \dots, s_l - 1, \quad (1.10)$$

where E is the $2n \times 2n$ identity matrix.

Using Lemma 1.1 and (1.10), we obtain

$$\mathfrak{x}_\zeta \nu^{0l} \gamma = \lambda_l^\zeta \nu^{0l} \gamma, \quad \mathfrak{x}_\zeta \nu^{\eta l} \gamma = \lambda_l^\zeta \nu^{\eta l} \gamma + \eta \nu^{\eta-1, l} \gamma \quad \text{for all } w \in \mathbb{C}^n, \quad \eta = 1, \dots, s_l - 1. \quad (1.11)$$

The following lemmas are needed for the sequel.

Lemma 1.2. *Let ν^{0l} be a common eigenvector of the matrices A_k corresponding to the eigenvalues λ_l^k , $k = 1, \dots, 2m$, respectively. Let $\nu^{\eta l}$, $\eta = 1, \dots, s_l - 1$ be generalized eigenvectors of the matrix A_ζ corresponding to the eigenvalue λ_l^ζ with elementary divisor of multiplicity s_l ($s_l \geq 2$). If the system (1.1.ζ) hasn't the first integrals*

$$F_{k\eta l}^\zeta: w \rightarrow \mathfrak{x}_k \Psi_{\eta l}^\zeta(w) \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m, \quad k \neq \zeta, \quad \eta = 1, \dots, s_l - 1, \quad (1.12)$$

then

$$\mathfrak{x}_\zeta \Psi_{\eta l}^\zeta(w) = \begin{cases} 1 & \text{for all } w \in \Omega, \quad \eta = 1, \\ 0 & \text{for all } w \in \Omega, \quad \eta = 2, \dots, s_l - 1, \end{cases} \quad (1.13)$$

$$\mathfrak{x}_k \Psi_{\eta l}^\zeta(w) = \mu_{\eta l}^{k\zeta} = \text{const} \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m, \quad k \neq \zeta, \quad \eta = 1, \dots, s_l - 1,$$

where $\Psi_{\eta l}^\zeta: \Omega \rightarrow \mathbb{C}$, $\eta = 1, \dots, s_l - 1$ is a solution to the system

$$\nu^{\eta l} \gamma = \sum_{\delta=1}^{\eta} \binom{\eta-1}{\delta-1} \Psi_{\delta l}^\zeta(w) \cdot \nu^{\eta-\delta, l} \gamma, \quad \eta = 1, \dots, s_l - 1, \quad \Omega \subset \{w: \nu^{0l} \gamma \neq 0\}. \quad (1.14)$$

Proof. The system (1.14) has the determinant $(\nu^{0l} \gamma)^{s_l-1}$. Therefore there exists the solution $\Psi_{\eta l}^\zeta$, $\eta = 1, \dots, s_l - 1$ on a domain $\Omega \subset \{w: \nu^{0l} \gamma \neq 0\}$ of the system (1.14).

The proof of the lemma is by induction on η .

For $\eta = 1$ and $\eta = 2$, the assertion (1.13) follows from (1.11).

Assume that (1.13) for $\eta = 1, \dots, \varepsilon - 1$ is true. Using (1.11) and (1.14), we get

$$\begin{aligned} \mathfrak{x}_\zeta \nu^{\varepsilon l} \gamma &= \lambda_l^\zeta \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \Psi_{\delta l}^\zeta(w) \nu^{\varepsilon-\delta, l} \gamma + (\varepsilon - 1) \sum_{\delta=1}^{\varepsilon-1} \binom{\varepsilon-2}{\delta-1} \Psi_{\delta l}^\zeta(w) \nu^{\varepsilon-\delta-1, l} \gamma + \\ &+ \nu^{\varepsilon-1, l} \gamma + \nu^{0l} \gamma \mathfrak{x}_\zeta \Psi_{\varepsilon l}^\zeta(w) \quad \text{for all } w \in \Omega. \end{aligned}$$

Combining (1.14) for $\eta = \varepsilon - 1$ and $\eta = \varepsilon$, (1.11) for $\eta = \varepsilon$, and $\nu^{0l} \gamma \neq 0$ in $\mathbb{C}^n$, we obtain $\mathfrak{x}_\zeta \Psi_{\varepsilon l}^\zeta(w) = 0$ for all $w \in \Omega$. So by the principle of mathematical induction, the statement (1.13) is true for every $\eta = 1, \dots, s_l - 1$ and $\zeta \in \{1, \dots, m\}$.

Taking into account (1.4) and (1.12), we have the statement (1.13) is true for $k \neq \zeta$. ■

Lemma 1.3 *Under the conditions of Lemma 1.2, we have*

$$\mathfrak{x}_k \nu^{\eta l} \gamma = \sum_{\delta=0}^{\eta} \binom{\eta}{\delta} \mu_{\delta l}^{k\zeta} \cdot \nu^{\eta-\delta, l} \gamma \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m, \quad \eta = 1, \dots, s_l - 1, \quad (1.15)$$

where $\mu_{0l}^{k\zeta} = \lambda_l^k$, $\mu_{\eta l}^{k\zeta} = \mathfrak{x}_k \Psi_{\eta l}^\zeta(w)$, $\eta = 1, \dots, s_l - 1$, $k = 1, \dots, 2m$.

Proof. The proof of Lemma 1.3 is by induction on η .

Let $\eta = 1$. Using (1.14), we get

$$\nu^{1l} \gamma = \Psi_{1l}^\zeta(w) \cdot \nu^{0l} \gamma \quad \text{for all } w \in \Omega, \quad \Omega \subset \{w: \nu^{0l} \gamma \neq 0\}. \quad (1.16)$$

Then

$$\mathfrak{x}_k \nu^{1l} \gamma = \mathfrak{x}_k \Psi_{1l}^\zeta(w) \cdot \nu^{0l} \gamma + \Psi_{1l}^\zeta(w) \cdot \mathfrak{x}_k \nu^{0l} \gamma \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m.$$

Taking into account Lemma 1.1, Lemma 1.2 and (1.16), we obtain

$$\mathfrak{r}_k \nu^{1l} \gamma = \mu_{1l}^{k\zeta} \cdot \nu^{0l} \gamma + \mu_{0l}^{k\zeta} \Psi_{1l}^\zeta(w) \cdot \nu^{0l} \gamma = \mu_{0l}^{k\zeta} \cdot \nu^{1l} \gamma + \mu_{1l}^{k\zeta} \cdot \nu^{0l} \gamma, \quad k = 1, \dots, 2m,$$

where $\mu_{0l}^{k\zeta} = \lambda_l^k$, $\mathfrak{r}_k \Psi_{1l}^\zeta(w) = \mu_{1l}^{k\zeta}$, $k = 1, \dots, 2m$. So (1.15) for $\eta = 1$ is true.

Suppose that the assertion of the lemma is valid for $\eta = 1, \dots, \varepsilon - 1$. From (1.14), we get

$$\mathfrak{r}_k \nu^{\varepsilon l} \gamma = \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \mathfrak{r}_k \Psi_{\delta l}^\zeta(w) \nu^{\varepsilon-\delta, l} \gamma + \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \Psi_{\delta l}^\zeta(w) \mathfrak{r}_k \nu^{\varepsilon-\delta, l} \gamma \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m.$$

By the induction hypothesis, we have

$$\begin{aligned} \mathfrak{r}_k \nu^{\varepsilon l} \gamma &= \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma + \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \Psi_{\delta l}^\zeta(w) \cdot \sum_{\varkappa=0}^{\varepsilon-\delta} \binom{\varepsilon-\delta}{\varkappa} \mu_{\varkappa l}^{k\zeta} \cdot \nu^{\varepsilon-\delta-\varkappa, l} \gamma = \\ &= \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma + \sum_{\alpha=0}^{\varepsilon-1} \binom{\varepsilon-1}{\alpha} \mu_{\alpha l}^{k\zeta} \cdot \sum_{\beta=1}^{\varepsilon-\alpha} \binom{\varepsilon-\alpha-1}{\beta-1} \Psi_{\beta l}^\zeta(w) \cdot \nu^{(\varepsilon-\alpha)-\beta, l} \gamma = \\ &= \sum_{\delta=1}^{\varepsilon} \binom{\varepsilon-1}{\delta-1} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma + \sum_{\alpha=0}^{\varepsilon-1} \binom{\varepsilon-1}{\alpha} \mu_{\alpha l}^{k\zeta} \cdot \nu^{\varepsilon-\alpha, l} \gamma = \\ &= \sum_{\delta=1}^{\varepsilon-1} \binom{\varepsilon-1}{\delta-1} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma + \mu_{\varepsilon l}^{k\zeta} \cdot \nu^{0l} \gamma + \mu_{0l}^{k\zeta} \cdot \nu^{\varepsilon l} \gamma + \sum_{\delta=1}^{\varepsilon-1} \binom{\varepsilon-1}{\delta} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma = \\ &= \mu_{0l}^{k\zeta} \cdot \nu^{\varepsilon l} \gamma + \sum_{\delta=1}^{\varepsilon-1} \binom{\varepsilon}{\delta} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma + \mu_{\varepsilon l}^{k\zeta} \cdot \nu^{0l} \gamma = \sum_{\delta=0}^{\varepsilon} \binom{\varepsilon}{\delta} \mu_{\delta l}^{k\zeta} \cdot \nu^{\varepsilon-\delta, l} \gamma, \quad k = 1, \dots, 2m. \end{aligned}$$

Thus by the principle of induction, the statement (1.15) is true for $\eta = 1, \dots, s_l - 1$. ■

Theorem 1.2. *Let the assumptions of Lemma 1.2 with $l = 1, \dots, r$ $\left(\sum_{l=1}^r s_l \geq m+1 \right)$ hold.*

Then the completely solvable system (1.1) has the autonomous first integral

$$F: w \rightarrow \prod_{\xi=1}^{\alpha} (\nu^{0\xi} \gamma)^{h_{0\xi}} \exp \sum_{q=1}^{\varepsilon_\xi} h_{q\xi} \Psi_{q\xi}^\zeta(w) \quad \text{for all } w \in \Omega, \quad \Omega \subset D(F), \quad (1.17)$$

where $\sum_{\xi=1}^{\alpha} \varepsilon_\xi = 2m - \alpha + 1$, $\varepsilon_\xi \leq s_\xi - 1$, $\xi = 1, \dots, \alpha$, $\alpha \leq r$, and $h_{q\xi}$, $q=0, \dots, \varepsilon_\xi$, $\xi=1, \dots, \alpha$ is a nontrivial solution to the linear homogeneous algebraic system of equations

$$\sum_{\xi=1}^{\alpha} (\lambda_\xi^k h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{k\zeta} h_{q\xi}) = 0, \quad k = 1, \dots, 2m.$$

Proof. The Lie derivative of (1.17) by virtue of (1.1) is equal to

$$\mathfrak{r}_k F(w) = \sum_{\xi=1}^{\alpha} (\lambda_\xi^k h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{k\zeta} h_{q\xi}) F(w) \quad \text{for all } w \in \Omega, \quad k = 1, \dots, 2m.$$

If $\sum_{\xi=1}^{\alpha} (\lambda_\xi^k h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{k\zeta} h_{q\xi}) = 0$, $k = 1, \dots, 2m$, then the $\mathbb{R}$ -differentiable function (1.17)

is an autonomous first integral of the completely solvable system (1.1). ■

As an example, the completely solvable $\mathbb{R}$ -linear system of equations in total differentials

$$\begin{aligned} dw_1 &= ((1+i)w_1 + iw_2 - \bar{w}_1 - \bar{w}_2)dz + (w_1 + iw_2 - \bar{w}_1 - \bar{w}_2)d\bar{z}, \\ dw_2 &= (w_2 + \bar{w}_1 + \bar{w}_2)dz + ((1-i)w_2 + \bar{w}_1 + \bar{w}_2)d\bar{z}, \\ dw_3 &= (-w_1 + w_2 + w_3 - i\bar{w}_2)dz + (-w_1 + w_2 + (1-i)w_3 - i\bar{w}_2)d\bar{z} \end{aligned} \quad (1.18)$$

has two eigenvalues: $\lambda_1^1 = 1+i$ with elementary divisor $(\lambda^1 - 1 - i)^3$ and $\lambda_2^1 = 1$ with elementary divisor $(\lambda^1 - 1)^3$. The $\lambda_1^1 = 1+i$ corresponding to the eigenvector $\nu^{01} = (1, 1, 0, 0, 0, 0)$ and to the generalized eigenvectors $\nu^{11} = (0, 0, 0, 0, 1, 0)$, $\nu^{21} = (0, 1, 0, 0, 0, 1)$. The eigenvalue $\lambda_2^1 = 1$ corresponding to the $\nu^{02} = (0, 0, 0, 1, 1, 0)$, $\nu^{12} = (0, 1, 0, 0, 0, 0)$, $\nu^{22} = (0, 0, 1, 0, 1, 0)$.

The scalar functions (see (1.14))

$$\Psi_{11}^1: w \rightarrow \frac{\bar{w}_2}{w_1 + w_2}, \quad \Psi_{21}^1: w \rightarrow \frac{(w_1 + w_2)(w_2 + \bar{w}_3) - \bar{w}_2^2}{(w_1 + w_2)^2} \quad \text{for all } w \in \Omega,$$

where $\Omega \subset \{w: w_1 + w_2 \neq 0\} \subset \mathbb{C}^3$. The $\mathbb{R}$ -differentiable first integrals (by Theorem 1.2)

$$F_1: w \rightarrow \Psi_{21}^1(w) \quad \text{for all } w \in \Omega \quad (1.19)$$

and

$$F_2: w \rightarrow \frac{\bar{w}_1 + \bar{w}_2}{w_1 + w_2} \exp\left(i \frac{\bar{w}_2}{w_1 + w_2}\right) \quad \text{for all } w \in \Omega \quad (1.20)$$

are an autonomous general integral of the system of equations in total differentials (1.18).

1.1.3. Nonautonomous $\mathbb{R}$ -differentiable first integrals

Theorem 1.3. Suppose ν is a common eigenvector of the matrices A_k corresponding to the eigenvalues λ^k , $k = 1, \dots, 2m$, respectively. Then the $\mathbb{R}$ -differentiable function

$$F: (z, w) \rightarrow (\nu\gamma) \exp\left(-\sum_{j=1}^m (\lambda^j z_j + \lambda^{m+j} \bar{z}_j)\right) \quad \text{for all } (z, w) \in \mathbb{C}^{m+n}$$

is a first integral of the system of equations in total differentials (1.1).

Proof. Using Lemma 1.1, we obtain $\mathfrak{X}_k F(z, w) = 0$ for all $(z, w) \in \mathbb{C}^{m+n}$, $k = 1, \dots, 2m$, where the linear nonautonomous differential operators

$$\mathfrak{X}_j(z, w) = \partial_{z_j} + \mathfrak{x}_j(w), \quad \mathfrak{X}_{m+j}(z, w) = \partial_{\bar{z}_j} + \mathfrak{x}_{m+j}(w) \quad \text{for all } (z, w) \in \mathbb{C}^{m+n}, j = 1, \dots, m. \blacksquare$$

Consider the system (1.8). Using the eigenvector $\nu^1 = (0, 1, 1, 1)$ corresponding to the eigenvalues $\lambda_1^1 = 1+i$ and $\lambda_2^1 = i$, we can build the first integral (by Theorem 1.3)

$$F: (z, w) \rightarrow (w_2 + \bar{w}_1 + \bar{w}_2) \exp(-(1+i)z - i\bar{z}) \quad \text{for all } (z, w) \in \mathbb{C}^3. \quad (1.21)$$

The first integrals (1.9) and (1.21) are a general integral of the system (1.8).

Theorem 1.4. Suppose the system (1.1) satisfies the conditions of Lemma 1.2. Then the completely solvable system (1.1) has the $\mathbb{R}$ -differentiable first integrals

$$F_\eta: (z, w) \rightarrow \Psi_{\eta l}^\zeta(w) - \sum_{j=1}^m (\mu_{\eta l}^{j\zeta} z_j + \mu_{\eta l}^{m+j, \zeta} \bar{z}_j) \quad \text{for all } (z, w) \in G, \quad \eta = 1, \dots, s_l - 1, \quad (1.22)$$

where a domain $G \subset \mathbb{C}^{m+n}$, the functions $\Psi_{\eta l}^\zeta$ are the solution to system (1.14), the numbers $\mu_{\eta l}^{k\zeta} = \mathfrak{x}_k \Psi_{\eta l}^\zeta(w)$, $\eta = 1, \dots, s_l - 1$, $k = 1, \dots, 2m$.

Proof. The Lie derivative of (1.22) by virtue of (1.1) is

$$\mathfrak{X}_k F_\eta(z, w) = -\mu_{\eta l}^{k\zeta} + \mathfrak{x}_k \Psi_{\eta l}^\zeta(w) \quad \text{for all } (z, w) \in G, \quad k = 1, \dots, 2m, \quad \eta = 1, \dots, s_l - 1.$$

Taking into account Lemma 1.2, we get the functions (1.22) are first integrals of (1.1). $\blacksquare$

For example, the system (1.18) has the numbers $\mu_{11}^{11} = 1$, $\mu_{11}^{21} = 1$, and the first integral

$$F: (z, w) \rightarrow \frac{\bar{w}_2}{w_1 + w_2} - z - \bar{z} \quad \text{for all } (z, w) \in \mathbb{C} \times \Omega \quad (\text{by Theorem 1.4}).$$

The $\mathbb{R}$ -differentiable first integrals (1.19), (1.20), and F are a general integral on a domain $\mathbb{C} \times \Omega$ of the system (1.18), where a domain $\Omega \subset \{w: w_1 + w_2 \neq 0\}$.

1.2. $\mathbb{R}$ -linear nonhomogeneous systems in total differentials

Let us consider a nonhomogeneous system of equations in total differentials

$$dw = \sum_{j=1}^m ((B_j \gamma + f_j(z)) dz_j + (B_{m+j} \gamma + f_{m+j}(z)) d\bar{z}_j) \quad (1.23)$$

corresponding to the $\mathbb{R}$ -linear homogeneous system (1.1), where the matrices $B_1, \dots, B_{2m}$ are transpose of $A_1, \dots, A_{2m}$, respectively, and the vector functions

$$f_k: z \rightarrow \text{colon}(f_{k1}(z), \dots, f_{kn}(z)) \quad \text{for all } z \in V, \quad k = 1, \dots, 2m$$

are continuously $\mathbb{R}$ -differentiable on a domain $V \subset \mathbb{C}^m$.

The Frobenius conditions for the total solvability of system of equations in total differentials (1.23) are the relations (1.4) and

$$\partial_{z_j} f^\zeta(z) + B_\zeta f^j(z) = \partial_{z_\zeta} f^j(z) + B_j f^\zeta(z) \quad \text{for all } z \in V, \quad j, \zeta = 1, \dots, m,$$

$$\partial_{\bar{z}_j} f^{m+\zeta}(z) + B_{m+\zeta} f^{m+j}(z) = \partial_{\bar{z}_\zeta} f^{m+j}(z) + B_{m+j} f^{m+\zeta}(z) \quad \text{for all } z \in V, \quad j, \zeta = 1, \dots, m,$$

$$\partial_{z_j} f^{m+\zeta}(z) + B_{m+\zeta} f^j(z) = \partial_{\bar{z}_\zeta} f^j(z) + B_j f^{m+\zeta}(z) \quad \text{for all } z \in V, \quad j, \zeta = 1, \dots, m,$$

where the vector functions

$$f^j: z \rightarrow \text{colon}(f_{j1}(z), \dots, f_{jn}(z), \bar{f}_{m+j,1}(z), \dots, \bar{f}_{m+j,n}(z)) \quad \text{for all } z \in V, \quad j = 1, \dots, m,$$

$$f^{m+j}: z \rightarrow \text{colon}(f_{m+j,1}(z), \dots, f_{m+j,n}(z), \bar{f}_{j1}(z), \dots, \bar{f}_{jn}(z)) \quad \text{for all } z \in V, \quad j = 1, \dots, m.$$

Theorem 1.5. *Let the assumptions of Lemma 1.2 hold. Then the completely solvable system of equations in total differentials (1.23) has the first integrals*

$$F_\eta: (z, w) \rightarrow \nu^{\eta l} \gamma \cdot \varphi(z) - \sum_{\tau=1}^{\eta} K_{\tau-1}^\eta(z) \cdot F_{\tau-1}(z, w) - C_\eta(z)$$

$$\text{for all } (z, w) \in \tilde{V} \times \Omega, \quad \eta = 0, \dots, s_l - 1, \quad \Omega \subset \{w: \nu^{0l} \gamma \neq 0\}, \quad \tilde{V} \subset V,$$

where the functions

$$\varphi: z \rightarrow \exp\left(-\sum_{k=1}^{2m} \mu_{0l}^{k\zeta} u_k\right) \quad \text{for all } z \in \tilde{V}, \quad u = (z_1, \dots, z_m, \bar{z}_1, \dots, \bar{z}_m),$$

$$K_{\tau-1}^\eta: z \rightarrow \int \sum_{k=1}^{2m} \left(\binom{\eta}{\tau-1} \mu_{\eta-\tau+1,l}^{k\zeta} + \sum_{\delta=1}^{\eta-\tau} \binom{\eta}{\delta} \mu_{\delta l}^{k\zeta} \cdot K_{\tau-1}^{\eta-\delta}(z) \right) du_k, \quad \tau = 1, \dots, \eta, \quad \eta = 1, \dots, s_l - 1,$$

$$C_\eta: z \rightarrow \int \sum_{k=1}^{2m} \left(\nu^{\eta l} f^k(z) \cdot \varphi(z) + \sum_{\tau=1}^{\eta} \binom{\eta}{\tau} \mu_{\tau l}^{k\zeta} \cdot C_{\eta-\tau}(z) \right) du_k \quad \text{for all } z \in \tilde{V}, \quad \eta = 0, \dots, s_l - 1,$$

the numbers $\mu_{0l}^{k\zeta} = \lambda_l^k$, $\mu_{\eta l}^{k\zeta} = \mathfrak{x}_k \Psi_{\eta l}^\zeta(w)$, $\eta = 1, \dots, s_l - 1$, $k = 1, \dots, 2m$.

2. First integrals of linear real systems in total differentials

2.1. Linear real homogeneous systems in total differentials

Consider an autonomous system of equations in total differentials

$$dx = A(x) dt, \quad (2.1)$$

where $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, $t = (t_1, \dots, t_m) \in \mathbb{R}^m$, the entries of the matrix $A(x) = \|a_{ij}(x)\|$ (with n rows and m columns) are linear homogeneous functions

$$a_{ij}: x \rightarrow \sum_{\xi=1}^n a_{ij\xi} x_\xi \quad \text{for all } x \in \mathbb{R}^n \quad (a_{ij\xi} \in \mathbb{R}, \xi = 1, \dots, n, j = 1, \dots, m, i = 1, \dots, n),$$

and $dx = \text{colon}(dx_1, \dots, dx_n)$ and $dt = \text{colon}(dt_1, \dots, dt_m)$ are vector columns.

Assume that this system is completely solvable. The Frobenius condition [5, p. 19] for the total solvability of system (2.1) in terms of the Poisson brackets is given by the relations

$$[\mathbf{p}_j(x), \mathbf{p}_\zeta(x)] = 0 \quad \text{for all } x \in \mathbb{R}^n, \quad j = 1, \dots, m, \quad \zeta = 1, \dots, m, \quad (2.2)$$

where the linear autonomous differential operators

$$\mathbf{p}_j(x) = \sum_{i=1}^n a_{ij}(x) \partial_{x_i} \quad \text{for all } x \in \mathbb{R}^n, \quad j = 1, \dots, m.$$

The Frobenius conditions (2.2) for system (2.1) are equivalent [60, p. 73]:

$$A_j A_\zeta = A_\zeta A_j, \quad j = 1, \dots, m, \quad \zeta = 1, \dots, m,$$

where $A_j = \|a_{\xi j}\|$, $j = 1, \dots, m$ are real $n \times n$ matrices.

2.1.1. Partial integrals. The complex-valued linear homogeneous function

$$p: x \rightarrow \sum_{\xi=1}^n b_\xi x_\xi \quad \text{for all } x \in \mathbb{R}^n \quad (b_\xi \in \mathbb{C}, \xi = 1, \dots, n)$$

is a *partial integral* of the system in total differentials (2.1) iff

$$\mathbf{p}_j p(x) = \lambda^j p(x) \quad \text{for all } x \in \mathbb{R}^n, \quad \lambda^j \in \mathbb{C}, \quad j = 1, \dots, m.$$

This system of identities is equivalent to the linear homogeneous system of equations $(A_j - \lambda^j E)b = 0$, $j = 1, \dots, m$, where E is the $n \times n$ identity matrix, $b = \text{colon}(b_1, \dots, b_n)$.

The proof of the following statement is similar to that of Lemma 1.1.

Lemma 2.1. *Suppose $\nu \in \mathbb{C}^n$ is a common eigenvector of the matrices A_j , $j = 1, \dots, m$. Then the linear function $p: x \rightarrow \nu x$ for all $x \in \mathbb{R}^n$ is a partial integral of the system (2.1).*

The following properties are needed for the sequel.

Property 2.1 ([11]). *Suppose $\nu = \nu^* + \tilde{\nu} i$ ($\nu^* = \text{Re } \nu$, $\tilde{\nu} = \text{Im } \nu$) is a common eigenvector of the matrices A_j corresponding to the eigenvalues $\lambda^j = \lambda^{*j} + \tilde{\lambda}^j i$ ($\lambda^{*j} = \text{Re } \lambda^j$, $\tilde{\lambda}^j = \text{Im } \lambda^j$), $j = 1, \dots, m$, respectively. Then the real-valued function*

$$P: x \rightarrow (\nu^* x)^2 + (\tilde{\nu} x)^2 \quad \text{for all } x \in \mathbb{R}^n$$

is a partial integral of the system of equations in total differentials (2.1) and

$$\mathbf{p}_j P(x) = 2 \lambda^{*j} P(x) \quad \text{for all } x \in \mathbb{R}^n, \quad j = 1, \dots, m.$$

Property 2.2 ([11]). Let $\nu = \overset{*}{\nu} + \tilde{\nu}i$ be a common eigenvector of the matrices A_j corresponding to the eigenvalues $\lambda^j = \overset{*}{\lambda}^j + \tilde{\lambda}^j i$, $j = 1, \dots, m$, respectively. Then

$$\mathfrak{p}_j \arctg \frac{\tilde{\nu}x}{\overset{*}{\nu}x} = \tilde{\lambda}^j \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m, \quad \mathcal{X} \subset \{x: \overset{*}{\nu}x \neq 0\}.$$

2.1.2. Autonomous first integrals. The proof of the following assertions is similar to those of Theorem 1.1 and Corollary 1.1.

Theorem 2.1. Suppose ν^k are real common eigenvectors of the matrices A_j corresponding to the eigenvalues λ_k^j , $j = 1, \dots, m$, $k = 1, \dots, m+1$, respectively. Then the system of equations in total differentials (2.1) has the autonomous first integral

$$F: x \rightarrow \prod_{k=1}^{m+1} |\nu^k x|^{h_k} \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset D(F),$$

where $h_1, \dots, h_{m+1}$ is a real nontrivial solution to the system $\sum_{k=1}^{m+1} \lambda_k^j h_k = 0$, $j = 1, \dots, m$.

Corollary 2.1. Let ν^k be real common eigenvectors of the matrices A_j corresponding to the eigenvalues λ_k^j , $j = 1, \dots, m$, $k = 1, \dots, m+1$, respectively. Then an autonomous first integral of the system of equations in total differentials (2.1) is the function

$$F_{12\dots m(m+1)}: x \rightarrow \prod_{k=1}^m |\nu^k x|^{-\Delta_k} |\nu^{m+1} x|^\Delta \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset D(F_{12\dots m(m+1)}),$$

where the determinants Δ_k , $k = 1, \dots, m$ are obtained by replacing the k -th column of the determinant $\Delta = |\lambda_k^j|$ by colon $(\lambda_{m+1}^1, \dots, \lambda_{m+1}^m)$, respectively.

As an example, the linear autonomous system of equations in total differentials

$$\begin{aligned} dx_1 &= -x_1 dt_2, & dx_2 &= 2(x_3 + x_4) dt_1 + x_2 dt_2, \\ dx_3 &= x_2 dt_1 + x_4 dt_2, & dx_4 &= x_2 dt_1 + x_3 dt_2 \end{aligned} \tag{2.3}$$

has the commuting matrices

$$A_1 = \begin{vmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 2 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{vmatrix} \quad \text{and} \quad A_2 = \begin{vmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{vmatrix}.$$

Therefore the system of equations in total differentials (2.3) is completely solvable.

The matrices A_1 and A_2 have the eigenvalues $\lambda_1^1 = -2$, $\lambda_2^1 = \lambda_3^1 = 0$, $\lambda_4^1 = 2$, and $\lambda_1^2 = 1$, $\lambda_2^2 = \lambda_3^2 = -1$, $\lambda_4^2 = 1$ corresponding to the eigenvectors $\nu^1 = (0, -1, 1, 1)$, $\nu^2 = (1, 0, 0, 0)$, $\nu^3 = (0, 0, 1, -1)$, $\nu^4 = (0, 1, 1, 1)$, respectively.

The determinants

$$\begin{aligned} \Delta &= \begin{vmatrix} -2 & 0 \\ 1 & -1 \end{vmatrix} = 2, & \Delta_{11} &= \begin{vmatrix} 0 & 0 \\ -1 & -1 \end{vmatrix} = 0, & \Delta_{21} &= \begin{vmatrix} -2 & 0 \\ 1 & -1 \end{vmatrix} = 2, \\ \Delta_{12} &= \begin{vmatrix} 2 & 0 \\ 1 & -1 \end{vmatrix} = -2, & \Delta_{22} &= \begin{vmatrix} -2 & 2 \\ 1 & 1 \end{vmatrix} = -4. \end{aligned}$$

By Corollary 2.1 we have that the functionally independent functions

$$F_{123}: x \rightarrow \frac{(x_3 - x_4)^2}{x_1^2} \quad \text{for all } x \in \mathcal{X} \tag{2.4}$$

and

$$F_{124}: x \rightarrow x_1^4 (x_2^2 - (x_3 + x_4)^2)^2 \quad \text{for all } x \in \mathbb{R}^4 \quad (2.5)$$

are first integrals of the system (2.3), where a domain $\mathcal{X} \subset \{x: x_1 \neq 0\}$.

Theorem 2.2. *Let $\nu^k = \nu^{*k} + \tilde{\nu}^k i$ (this set hasn't complex conjugate vectors) be common complex eigenvectors of the matrices A_j corresponding to the eigenvalues $\lambda_k^j = \lambda_k^{*j} + \tilde{\lambda}_k^j i$ $j = 1, \dots, m$, $k = 1, \dots, s$, $s \leq (m+1)/2$, respectively. Let ν^θ be common real eigenvectors of A_j corresponding to the eigenvalues λ_θ^j , $j = 1, \dots, m$, $\theta = s+1, \dots, m+1-s$. Then the system of equations in total differentials (2.1) has the autonomous first integral*

$$F: x \rightarrow \prod_{k=1}^s (P_k(x))^{\tilde{h}_k} \exp\left(-2\tilde{h}_k \varphi_k(x)\right) \prod_{\theta=s+1}^{m+1-s} |\nu^\theta x|^{h_\theta} \quad \text{for all } x \in \mathcal{X}, \quad (2.6)$$

where a domain $\mathcal{X} \subset D(F)$, the functions

$$P_k: x \rightarrow (\nu^{*k} x)^2 + (\tilde{\nu}^k x)^2 \quad \text{for all } x \in \mathbb{R}^n, \quad \varphi_k: x \rightarrow \arctg \frac{\tilde{\nu}^k x}{\nu^{*k} x} \quad \text{for all } x \in \mathcal{X}, \quad k = 1, \dots, s,$$

and $\tilde{h}_k, \tilde{h}_k, k = 1, \dots, s$, $h_\theta, \theta = s+1, \dots, m+1-s$ is a real nontrivial solution to

$$2 \sum_{k=1}^s (\lambda_k^{*j} \tilde{h}_k - \tilde{\lambda}_k^j \tilde{h}_k) + \sum_{\theta=s+1}^{m+1-s} \lambda_\theta^j h_\theta = 0, \quad j = 1, \dots, m.$$

Proof. Taking into account Property 2.1 and Lemma 2.1, we obtain

$$\begin{aligned} \mathfrak{p}_j((\nu^{*k} x)^2 + (\tilde{\nu}^k x)^2) &= 2 \lambda_k^{*j} ((\nu^{*k} x)^2 + (\tilde{\nu}^k x)^2), \quad j = 1, \dots, m, \quad k = 1, \dots, s, \\ \mathfrak{p}_j \nu^\theta x &= \lambda_\theta^j \nu^\theta x \quad \text{for all } x \in \mathbb{R}^n, \quad j = 1, \dots, m, \quad \theta = s+1, \dots, m+1-s. \end{aligned} \quad (2.7)$$

The Lie derivative of (2.6) by virtue of (2.1) is equal to

$$\begin{aligned} \mathfrak{p}_j F(x) &= \left(\prod_{k=1}^s (P_k(x))^{\tilde{h}_k-1} \exp\left(-2\tilde{h}_k \varphi_k(x)\right) \sum_{k=1}^s \tilde{h}_k \prod_{l=1, l \neq k}^s P_l(x) \cdot \mathfrak{p}_j P_k(x) + \right. \\ &\quad \left. + \prod_{k=1}^s (P_k(x))^{\tilde{h}_k} \exp\left(-2\tilde{h}_k \varphi_k(x)\right) \sum_{k=1}^s \mathfrak{p}_j \left(-2\tilde{h}_k \varphi_k(x)\right) \right) \prod_{\theta=s+1}^{m+1-s} |\nu^\theta x|^{h_\theta} + \\ &\quad + \prod_{k=1}^s (P_k(x))^{\tilde{h}_k} \exp\left(-2\tilde{h}_k \varphi_k(x)\right) \prod_{\theta=s+1}^{m+1-s} |\nu^\theta x|^{h_\theta-1} \cdot \\ &\quad \cdot \sum_{\theta=s+1}^{m+1-s} \operatorname{sgn}(\nu^\theta x) h_\theta \prod_{l=s+1, l \neq \theta}^{m+1-s} |\nu^l x| \cdot \mathfrak{p}_j(\nu^\theta x) \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m. \end{aligned}$$

Using Property 2.2 and (2.7), we get

$$\mathfrak{p}_j F(x) = \left(\sum_{k=1}^s 2(\lambda_k^{*j} \tilde{h}_k - \tilde{\lambda}_k^j \tilde{h}_k) + \sum_{\theta=s+1}^{m+1-s} \lambda_\theta^j h_\theta \right) F(x) \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m.$$

If $2 \sum_{k=1}^s (\lambda_k^{*j} \tilde{h}_k - \tilde{\lambda}_k^j \tilde{h}_k) + \sum_{\theta=s+1}^{m+1-s} \lambda_\theta^j h_\theta = 0$, $j = 1, \dots, m$, then the function (2.6) is an

autonomous first integral of the system of equations in total differentials (2.1). ■

For example, the completely solvable linear system of total differential equations

$$dx_1 = x_1 dt_1 + x_2 dt_2, \quad dx_2 = x_2 dt_1 - x_1 dt_2, \quad dx_3 = x_3 dt_1 - x_3 dt_2 \quad (2.8)$$

has the eigenvalues $\lambda_1^1 = \lambda_2^1 = \lambda_3^1 = 1$, $\lambda_1^2 = -i$, $\lambda_2^2 = i$, $\lambda_3^2 = -1$ corresponding to the eigenvectors $\nu^1 = (1, i, 0)$, $\nu^2 = (1, -i, 0)$, $\nu^3 = (0, 0, 1)$, respectively. According to Theorem 2.2, we can construct the autonomous first integral of the system (2.8):

$$F: x \rightarrow \frac{x_1^2 + x_2^2}{x_3^2} \exp\left(2 \operatorname{arctg} \frac{x_2}{x_1}\right) \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 \neq 0, x_3 \neq 0\}. \quad (2.9)$$

Theorem 2.3. Suppose $\nu^\tau = \nu^{\tau*} + \tilde{\nu}^\tau i$, $\nu^{s+\tau} = \nu^{s+\tau*} - \tilde{\nu}^\tau i$, $\tau = 1, \dots, s$, $s \leq m/2$, and $\nu^{2s+1} = \nu^{2s+1*} + \tilde{\nu}^{2s+1} i$ are common complex eigenvectors of the matrices A_j corresponding to the eigenvalues $\lambda_\tau^j = \lambda_\tau^{j*} + \tilde{\lambda}_\tau^j i$, $\lambda_{s+\tau}^j = \lambda_\tau^{j*} - \tilde{\lambda}_\tau^j i$, $\tau = 1, \dots, s$, $\lambda_{2s+1}^j = \lambda_{2s+1}^{j*} + \tilde{\lambda}_{2s+1}^j i$, $j = 1, \dots, m$, respectively. Let ν^θ be common real eigenvectors of A_j corresponding to the eigenvalues λ_θ^j , $j = 1, \dots, m$, $\theta = 2s+2, \dots, m+1$, respectively. Then the system of equations in total differentials (2.1) has the autonomous first integrals

$$F_1: x \rightarrow \prod_{k=1}^s (P_k(x))^{\tilde{h}_k + \tilde{h}_{s+k}} \exp\left(-2(\tilde{h}_k - \tilde{h}_{s+k})\varphi_k(x)\right) \cdot (P_{2s+1}(x))^{\tilde{h}_{2s+1}} \exp\left(-2\tilde{h}_{2s+1}\varphi_{2s+1}(x)\right) \prod_{\theta=2s+2}^{m+1} (\nu^\theta x)^{2\tilde{h}_\theta} \quad \text{for all } x \in \mathcal{X} \quad (2.10)$$

and

$$F_2: x \rightarrow \prod_{k=1}^s (P_k(x))^{\tilde{h}_k + \tilde{h}_{s+k}} \exp\left(2(\tilde{h}_k - \tilde{h}_{s+k})\varphi_k(x)\right) \cdot (P_{2s+1}(x))^{\tilde{h}_{2s+1}} \exp\left(2\tilde{h}_{2s+1}\varphi_{2s+1}(x)\right) \prod_{\theta=2s+2}^{m+1} (\nu^\theta x)^{2\tilde{h}_\theta} \quad \text{for all } x \in \mathcal{X}, \quad (2.11)$$

where a domain $\mathcal{X} \subset D(F_1) \cap D(F_2)$, the functions

$$P_k: x \rightarrow (\nu^k x)^2 + (\tilde{\nu}^k x)^2 \quad \text{for all } x \in \mathbb{R}^n, \quad k = 1, \dots, s, 2s+1,$$

$$\varphi_k: x \rightarrow \operatorname{arctg} \frac{\tilde{\nu}^k x}{\nu^k x} \quad \text{for all } x \in \mathcal{X}, \quad k = 1, \dots, s, 2s+1,$$

and $h_k = h_k^* + \tilde{h}_k i$, $k = 1, \dots, m+1$ is a nontrivial solution to $\sum_{k=1}^{m+1} \lambda_k^j h_k = 0$, $j = 1, \dots, m$.

Proof. We form two complex-valued functions

$$F^*: x \rightarrow \prod_{k=1}^{2s} (\nu^k x)^{h_k} (\nu^{2s+1} x)^{h_{2s+1}} \prod_{\theta=2s+2}^{m+1} (\nu^\theta x)^{h_\theta} \quad \text{for all } x \in \mathcal{X}$$

and

$$F^{**}: x \rightarrow \prod_{k=1}^{2s} (\nu^k x)^{l_k} (\overline{\nu^{2s+1} x})^{l_{2s+1}} \prod_{\theta=2s+2}^{m+1} (\nu^\theta x)^{l_\theta} \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \mathbb{R}^n$ and h_k , l_k , $k = 1, \dots, m+1$ are complex numbers.

Using Lemma 2.1, we get

$$\mathfrak{p}_j^* F(x) = \sum_{k=1}^{m+1} \lambda_k^j h_k^* F(x) \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m,$$

$$\mathfrak{p}_j^{**} F(x) = \left(\sum_{k=1}^{2s} \lambda_k^j l_k + \overline{\lambda_{2s+1}^j} l_{2s+1} + \sum_{\theta=2s+2}^{m+1} \lambda_\theta^j l_\theta \right)^{**} F(x) \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m.$$

Let $h_k = h_k^* + \tilde{h}_k i$, $k=1, \dots, m+1$ be a nontrivial solution to $\sum_{k=1}^{m+1} \lambda_k^j h_k = 0$, $j=1, \dots, m$. Then $l_k = h_{s+k}^* - \tilde{h}_{s+k} i$, $l_{s+k} = h_k^* - \tilde{h}_k i$, $k=1, \dots, s$, $l_{2s+1} = h_{2s+1}^* - \tilde{h}_{2s+1} i$, $l_\theta = h_\theta^* - \tilde{h}_\theta i$, $\theta = 2s+2, \dots, m+1$ is a solution to $\sum_{k=1}^{2s} \lambda_k^j l_k + \overline{\lambda_{2s+1}^j} l_{2s+1} + \sum_{\theta=2s+2}^{m+1} \lambda_\theta^j l_\theta = 0$, $j=1, \dots, m$ and the functions $F^*: \mathcal{X} \rightarrow \mathbb{C}$, $F^{**}: \mathcal{X} \rightarrow \mathbb{C}$ are first integrals of the system (2.1).

Since $F_1 = F^* F^{**}$ and $F_2 = (F^*/F^*)^i$, we see that the functions (2.10) and (2.11) are autonomous first integrals of the system of equations in total differentials (2.1). ■

In particular, the completely solvable linear system of total differential equations

$$\begin{aligned} dx_1 &= x_1 dt_1 + x_3 dt_2, & dx_2 &= -x_2 dt_1 + x_4 dt_2, \\ dx_3 &= x_3 dt_1 - x_1 dt_2, & dx_4 &= -x_4 dt_1 - x_2 dt_2 \end{aligned} \quad (2.12)$$

has the eigenvalues $\lambda_1^1 = \lambda_2^1 = -1$, $\lambda_3^1 = \lambda_4^1 = 1$, $\lambda_1^2 = \lambda_3^2 = -i$, $\lambda_2^2 = \lambda_4^2 = i$ corresponding to the linearly independent eigenvectors $\nu^1 = (0, -i, 0, 1)$, $\nu^2 = (0, i, 0, 1)$, $\nu^3 = (-i, 0, 1, 0)$, and $\nu^4 = (i, 0, 1, 0)$, respectively. The functions (by Theorem 2.3)

$$F_1: x \rightarrow \frac{x_1 x_2 + x_3 x_4}{x_1 x_4 - x_2 x_3} \quad \text{for all } x \in \mathcal{X}, \quad F_2: x \rightarrow (x_1^2 + x_3^2)(x_2^2 + x_4^2) \quad \text{for all } x \in \mathbb{R}^4 \quad (2.13)$$

are first integrals of the system (2.12), where a domain $\mathcal{X} \subset \{x: x_1 x_4 - x_2 x_3 \neq 0\} \subset \mathbb{R}^4$.

From the entire set of ordinary differential systems induced by the completely solvable system of equations in total differentials (2.1), we extract system

$$dx = A^\zeta(x) dt_\zeta, \quad A^\zeta(x) = \text{colon}(a_{1\zeta}(x), \dots, a_{n\zeta}(x)) \quad \text{for all } x \in \mathbb{R}^n, \quad (2.1.\zeta)$$

such that the matrix A_ζ has the smallest number of elementary divisors.

Definition 2.1. Let ν^{0l} be an eigenvector of the matrix A_ζ corresponding to the eigenvalue λ_l^ζ with elementary divisor of multiplicity s_l . A non-zero vector $\nu^{\theta l} \in \mathbb{C}^n$ is called a **generalized eigenvector of order θ** for λ_l^ζ if and only if

$$(A_\zeta - \lambda_l^\zeta E) \nu^{\theta l} = \theta \cdot \nu^{\theta-1, l}, \quad \theta = 1, \dots, s_l - 1, \quad (2.14)$$

where E is the $n \times n$ identity matrix.

Using Lemma 2.1 and (2.14), we obtain

$$\mathfrak{p}_\zeta \nu^{0l} x = \lambda_l^\zeta \nu^{0l} x, \quad \mathfrak{p}_\zeta \nu^{\theta l} x = \lambda_l^\zeta \nu^{\theta l} x + \theta \nu^{\theta-1, l} x \quad \text{for all } x \in \mathbb{R}^n, \quad \theta = 1, \dots, s_l - 1. \quad (2.15)$$

The following lemmas are needed for the sequel.

Lemma 2.2. Let ν^{0l} be a common eigenvector of the matrices A_j corresponding to the eigenvalues λ_l^j , $j = 1, \dots, m$, respectively. Let $\nu^{\theta l}$, $\theta = 1, \dots, s_l - 1$ be generalized eigenvectors of the matrix A_ζ corresponding to the eigenvalue λ_l^ζ with elementary divisor of multiplicity s_l ($s_l \geq 2$). If the system (2.1.\zeta) hasn't the first integrals

$$F_{j\theta l}^\zeta: x \rightarrow \mathfrak{p}_j v_{\theta l}^\zeta(x) \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m, \quad j \neq \zeta, \quad \theta = 1, \dots, s_l - 1, \quad (2.16)$$

then

$$\mathfrak{p}_\zeta v_{\theta l}^\zeta(x) = \begin{cases} 1 & \text{for all } x \in \mathcal{X}, \quad \theta = 1, \\ 0 & \text{for all } x \in \mathcal{X} \quad \theta = 2, \dots, s_l - 1, \end{cases}$$

$$\mathfrak{p}_j v_{\theta l}^\zeta(x) = \mu_{\theta l}^{j\zeta} = \text{const} \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m, \quad j \neq \zeta, \quad \theta = 1, \dots, s_l - 1,$$

where $v_{\theta l}^\zeta: \mathcal{X} \rightarrow \mathbb{R}$, $\theta = 1, \dots, s_l - 1$ is a solution to the system

$$\nu^{\theta l} x = \sum_{\delta=1}^{\theta} \binom{\theta-1}{\delta-1} v_{\delta l}^\zeta(x) \cdot \nu^{\theta-\delta, l} x, \quad \theta = 1, \dots, s_l - 1, \quad \mathcal{X} \subset \{x: \nu^{0l} x \neq 0\}. \quad (2.17)$$

Lemma 2.3 Under the conditions of Lemma 2.2, we have

$$\mathfrak{p}_j \nu^{\theta l} x = \sum_{\delta=0}^{\theta} \binom{\theta}{\delta} \mu_{\delta l}^{j\zeta} \cdot \nu^{\theta-\delta, l} x \quad \text{for all } x \in \mathcal{X}, \quad j = 1, \dots, m, \quad \theta = 1, \dots, s_l - 1,$$

where $\mu_{0l}^{j\zeta} = \lambda_l^j$, $\mu_{\theta l}^{j\zeta} = \mathfrak{p}_j v_{\theta l}^\zeta(x)$, $\theta = 1, \dots, s_l - 1$, $j = 1, \dots, m$.

Theorem 2.4. Let the assumptions of Lemma 2.2 with $l = 1, \dots, r$ $\left(\sum_{l=1}^r s_l \geq m+1 \right)$ hold.

Then the completely solvable system (2.1) has the autonomous first integral

$$F: x \rightarrow \prod_{\xi=1}^k (\nu^{0\xi} x)^{h_{0\xi}} \exp \sum_{q=1}^{\varepsilon_\xi} h_{q\xi} v_{q\xi}^\zeta(x) \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset D(F),$$

where $\sum_{\xi=1}^k \varepsilon_\xi = m - k + 1$, $\varepsilon_\xi \leq s_\xi - 1$, $\xi = 1, \dots, k$, $k \leq r$, and $h_{q\xi}$, $q = 0, \dots, \varepsilon_\xi$, $\xi = 1, \dots, k$ is a nontrivial solution to the linear homogeneous algebraic system of equations

$$\sum_{\xi=1}^k (\lambda_\xi^j h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{j\zeta} h_{q\xi}) = 0, \quad j = 1, \dots, m.$$

The completely solvable linear homogeneous system of total differential equations

$$\begin{aligned} dx_1 &= x_2 dt_1 + (2x_1 - x_3) dt_2, \\ dx_2 &= (2x_2 - x_3 - x_4) dt_1 + (-x_1 + 2x_2 + x_4) dt_2, \\ dx_3 &= (x_1 - x_4) dt_1 + (-x_1 + 3x_3 + x_4) dt_2, \\ dx_4 &= (-x_1 + 2x_3 + 2x_4) dt_1 + (x_2 - 3x_3 + x_4) dt_2 \end{aligned} \quad (2.18)$$

has the eigenvalue $\lambda_1^1 = 1$ with elementary divisor $(\lambda^1 - 1)^4$ corresponding to the eigenvector $\nu^0 = (-1, 1, -1, 0)$ and to the generalized eigenvectors $\nu^1 = (1, 0, -1, -1)$, $\nu^2 = (1, -1, 3, 0)$, $\nu^3 = (-3, 0, 9, 9)$. The functions (see (2.17))

$$\begin{aligned} v_{11}^1: x &\rightarrow \frac{x_1 - x_3 - x_4}{-x_1 + x_2 - x_3} \quad \text{for all } x \in \mathcal{X}, \\ v_{21}^1: x &\rightarrow \frac{(-x_1 + x_2 - x_3)(x_1 - x_2 + 3x_3) - (x_1 - x_3 - x_4)^2}{(-x_1 + x_2 - x_3)^2} \quad \text{for all } x \in \mathcal{X}, \end{aligned} \quad (2.19)$$

$$\begin{aligned} v_{31}^1: x &\rightarrow \frac{1}{(-x_1 + x_2 - x_3)^3} \left((-3x_1 + 9x_3 + 9x_4)(-x_1 + x_2 - x_3)^2 - \right. \\ &\quad \left. - 3(-x_1 + x_2 - x_3)(x_1 - x_3 - x_4)(x_1 - x_2 + 3x_3) + 2(x_1 - x_3 - x_4)^3 \right) \quad \text{for all } x \in \mathcal{X}, \end{aligned}$$

where a domain $\mathcal{X} \subset \{x: x_1 - x_2 + x_3 \neq 0\}$.

Autonomous first integrals of the system (2.18) are the functions (by Theorem 2.4)

$$F_1: x \rightarrow v_{21}^1(x), \quad F_2: x \rightarrow (-x_1 + x_2 - x_3)^2 \exp(-2v_{11}^1(x) - v_{31}^1(x)) \quad \text{for all } x \in \mathcal{X}. \quad (2.20)$$

In the complex case, we shall have two logical possibilities:

1. Any function from the set $V = \{\nu^{0\xi}x, v_{q\xi}^\zeta(x) : q=1, \dots, \varepsilon_\xi, \xi=1, \dots, k, \sum_{\xi=1}^k \varepsilon_\xi = m-k+1\}$

has the complex conjugate function in the set V .

2. At least one function from the set V has not the complex conjugate function in the V .

Case 1. The completely solvable system (2.1) has the autonomous first integral

$$F: x \rightarrow \prod_{\xi=1}^{k_1} ((\nu^{0\xi}x)^2 + (\tilde{\nu}^{0\xi}x)^2)^{h_{0\xi}} \exp\left(-2\tilde{h}_{0\xi} \arctg \frac{\tilde{\nu}^{0\xi}x}{\nu^{0\xi}x} + \right. \\ \left. + 2 \sum_{q=1}^{\varepsilon_\xi} (h_{q\xi}^* v_{q\xi}^\zeta(x) - \tilde{h}_{q\xi} \tilde{v}_{q\xi}^\zeta(x))\right) \prod_{\theta=1}^{k_2} |\nu^{0\theta}x|^{h_{0\theta}} \exp \sum_{q=1}^{\varepsilon_\theta} h_{q\theta} v_{q\theta}^\zeta(x) \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset D(F),$$

where $h_{q\xi}^*, \tilde{h}_{q\xi}, q=0, \dots, \varepsilon_\xi, \xi=1, \dots, k_1, h_{q\theta}, q=0, \dots, \varepsilon_\theta, \theta=1, \dots, k_2$ is a real non-trivial solution to the linear homogeneous algebraic system of equations

$$2 \sum_{\xi=1}^{k_1} ((\lambda_\xi^j h_{0\xi}^* - \tilde{\lambda}_\xi^j \tilde{h}_{0\xi}) + \sum_{q=1}^{\varepsilon_\xi} (\mu_{q\xi}^{j\zeta} h_{q\xi}^* - \tilde{\mu}_{q\xi}^{j\zeta} \tilde{h}_{q\xi})) + \sum_{\theta=1}^{k_2} (\lambda_\theta^j h_{0\theta} + \sum_{q=1}^{\varepsilon_\theta} \mu_{q\theta}^{j\zeta} h_{q\theta}) = 0, \quad j=1, \dots, m.$$

Here $\nu^{0\xi} = \nu^{0\xi} + \tilde{\nu}^{0\xi}i$ are complex common eigenvectors of the matrices A_j corresponding to the eigenvalues $\lambda_\xi^j = \lambda_\xi^j + \tilde{\lambda}_\xi^j i, j=1, \dots, m, \xi=1, \dots, k_1$, respectively; $\nu^{0\theta}$ are real common eigenvectors of A_j corresponding to the eigenvalues $\lambda_\theta^j, j=1, \dots, m, \theta=1, \dots, k_2$, respectively; the functions $v_{q\xi}^\zeta = v_{q\xi}^\zeta + \tilde{v}_{q\xi}^\zeta i, v_{q\theta}^\zeta$ is the solution to the system (2.17); the numbers

$$\mu_{q\xi}^{j\zeta} = \mathbf{p}_j \operatorname{Re} v_{q\xi}^\zeta = \mathbf{p}_j v_{q\xi}^{\zeta*}, \quad \tilde{\mu}_{q\xi}^{j\zeta} = \mathbf{p}_j \operatorname{Im} v_{q\xi}^\zeta = \mathbf{p}_j \tilde{v}_{q\xi}^\zeta, \quad q=1, \dots, \varepsilon_\xi, \quad \xi=1, \dots, k_1,$$

$$\mu_{q\theta}^{j\zeta} = \mathbf{p}_j v_{q\theta}^\zeta, \quad q=1, \dots, \varepsilon_\theta, \quad \theta=1, \dots, k_2, \quad j=1, \dots, m;$$

the numbers $\varepsilon_\xi, \varepsilon_\theta$ such that $2 \sum_{\xi=1}^{k_1} \varepsilon_\xi + \sum_{\theta=1}^{k_2} \varepsilon_\theta = m - 2k_1 - k_2 + 1$ with $2k_1 + k_2 \leq r, \varepsilon_\xi \leq s_\xi - 1, \xi=1, \dots, k_1, \varepsilon_\theta \leq s_\theta - 1, \theta=1, \dots, k_2$, where k_1 is a number of complex common eigenvectors (this set hasn't complex conjugate vectors) of the matrices A_j and k_2 is a number of real common eigenvectors of the matrices $A_j, j=1, \dots, m$.

For example, the completely solvable system of total differential equations

$$\begin{aligned} dx_1 &= (3, -4, 4, 1, 0, 2)x dt_1 + (0, -4, 2, 1, -1, 1)x dt_2 - (3, -2, 4, 3, 0, 2)x dt_3, \\ dx_2 &= -(1, -3, 3, 0, 2, 3)x dt_1 + (1, 3, 0, 0, 1, -1)x dt_2 + (2, -3, 3, 3, -1, 2)x dt_3, \\ dx_3 &= -(3, -5, 5, 1, 2, 4)x dt_1 - (0, -6, 2, 1, -2, 1)x dt_2 + (3, -3, 5, 4, 0, 2)x dt_3, \\ dx_4 &= (3, -6, 4, 4, -1, 5)x dt_1 + (2, -6, 2, 3, -4, 2)x dt_2 - (3, -2, 6, 4, 1, 1)x dt_3, \\ dx_5 &= (5, -5, 8, 3, 3, 6)x dt_1 + (1, -6, 3, 2, -2, 2)x dt_2 - (3, -3, 6, 4, 1, 2)x dt_3, \\ dx_6 &= -(2, -5, 4, 3, -1, 2)x dt_1 - (2, -4, 3, 3, -2, 2)x dt_2 + (2, -1, 4, 2, 1, 0)x dt_3 \end{aligned} \quad (2.21)$$

has the eigenvalue $\lambda_1^1 = 1 + 2i$ with elementary divisor $(\lambda^1 - 1 - 2i)^3$ corresponding to the eigenvector $\nu^0 = (1, 0, 1 + i, 1, i, 1)$ and to the generalized eigenvectors $\nu^1 = (1, 1 + i, 0, 0, i, i)$,

$\nu^2 = (2 + 2i, 0, 2 + 2i, 0, 2i, 2i)$. The scalar functions of the vector argument

$$\begin{aligned}
v_{11}^* &: x \rightarrow ((x_1 + x_2)(x_1 + x_3 + x_4 + x_6) + (x_3 + x_5)(x_2 + x_5 + x_6))/P(x), \\
\tilde{v}_{11}^1 &: x \rightarrow ((x_1 + x_3 + x_4 + x_6)(x_2 + x_5 + x_6) - (x_1 + x_2)(x_3 + x_5))/P(x), \\
v_{21}^* &: x \rightarrow (((x_1 + x_3 + x_4 + x_6)(x_2 + x_5 + x_6) - (x_1 + x_2)(x_3 + x_5))^2 + \\
&+ 2P(x)((x_1 + x_3)(x_1 + x_3 + x_4 + x_6) + (x_3 + x_5)(x_1 + x_3 + x_5 + x_6)) - \\
&- ((x_1 + x_2)(x_1 + x_3 + x_4 + x_6) + (x_3 + x_5)(x_2 + x_5 + x_6))^2)/P(x), \\
\tilde{v}_{21}^1 &: x \rightarrow 2(P(x)((x_1 + x_3 + x_4 + x_6)(x_1 + x_3 + x_5 + x_6) - (x_1 + x_3)(x_3 + x_5)) + \\
&+ ((x_3 + x_5)(x_1 + x_2) - (x_1 + x_3 + x_4 + x_6)(x_2 + x_5 + x_6)) \cdot \\
&\cdot ((x_1 + x_2)(x_1 + x_3 + x_4 + x_6) + (x_3 + x_5)(x_2 + x_5 + x_6)))/P(x) \quad \text{for all } x \in \mathcal{X},
\end{aligned} \tag{2.22}$$

where a domain $\mathcal{X} \subset \{x: x_1 + x_3 + x_4 + x_6 \neq 0\}$, the polynomial

$$P: x \rightarrow (x_1 + x_3 + x_4 + x_6)^2 + (x_3 + x_5)^2 \quad \text{for all } x \in \mathbb{R}^6. \tag{2.23}$$

Autonomous first integrals of the system (2.21) are the functions

$$F_1: x \rightarrow P(x) \exp(-4\varphi(x) + 6v_{11}^*(x) + 2\tilde{v}_{11}^1(x)) \quad \text{for all } x \in \mathcal{X}, \tag{2.24}$$

$$F_2: x \rightarrow P^2(x) \exp(-2\varphi(x) + v_{21}^*(x) - \tilde{v}_{21}^1(x)) \quad \text{for all } x \in \mathcal{X}, \tag{2.25}$$

$$F_3: x \rightarrow 2\tilde{v}_{11}^1(x) - 2v_{21}^*(x) - \tilde{v}_{21}^1(x) \quad \text{for all } x \in \mathcal{X}, \tag{2.26}$$

where

$$\varphi: x \rightarrow \arctg \frac{x_3 + x_5}{x_1 + x_3 + x_4 + x_6} \quad \text{for all } x \in \mathcal{X}. \tag{2.27}$$

Case 2A. A common complex eigenvector of the matrices $A_j, j=1, \dots, m$ hasn't the complex conjugate vector. First integrals of the completely solvable system (2.1) are the functions

$$\begin{aligned}
F_1: x &\rightarrow \prod_{\xi=1}^{k_1} (P_\xi(x))^{h_{0\xi} + h_{0,(k_1+\xi)}} \exp\left(-2(\tilde{h}_{0\xi} - \tilde{h}_{0,(k_1+\xi)})\varphi_\xi(x) + \right. \\
&+ 2 \sum_{q=1}^{\varepsilon_\xi} \left((h_{q\xi}^* + h_{q,(k_1+\xi)}^*) v_{q\xi}^\zeta(x) + (\tilde{h}_{q,(k_1+\xi)} - \tilde{h}_{q\xi}) \tilde{v}_{q\xi}^\zeta(x) \right) (P_{2k_1+1}(x))^{h_{0,(2k_1+1)}} \cdot \\
&\cdot \exp\left(-2\tilde{h}_{0,(2k_1+1)} \varphi_{2k_1+1}(x)\right) \cdot \prod_{\theta=1}^{k_2} (\nu^{0\theta} x)^{2h_{0\theta}} \exp\left(2 \sum_{q=1}^{\varepsilon_\theta} h_{q\theta}^* v_{q\theta}^\zeta(x)\right) \quad \text{for all } x \in \mathcal{X}, \\
F_2: x &\rightarrow \prod_{\xi=1}^{k_1} (P_\xi(x))^{\tilde{h}_{0\xi} + \tilde{h}_{0,(k_1+\xi)}} \exp\left(2(h_{0\xi}^* - h_{0,(k_1+\xi)}^*) \varphi_\xi(x) + \right. \\
&+ 2 \sum_{q=1}^{\varepsilon_\xi} \left((\tilde{h}_{q\xi} + \tilde{h}_{q,(k_1+\xi)}) v_{q\xi}^\zeta(x) + (h_{q\xi}^* - h_{q,(k_1+\xi)}^*) \tilde{v}_{q\xi}^\zeta(x) \right) (P_{2k_1+1}(x))^{\tilde{h}_{0,(2k_1+1)}} \cdot \\
&\cdot \exp\left(2\tilde{h}_{0,(2k_1+1)} \varphi_{2k_1+1}(x)\right) \cdot \prod_{\theta=1}^{k_2} (\nu^{0\theta} x)^{2\tilde{h}_{0\theta}} \exp\left(2 \sum_{q=1}^{\varepsilon_\theta} \tilde{h}_{q\theta} v_{q\theta}^\zeta(x)\right) \quad \text{for all } x \in \mathcal{X},
\end{aligned}$$

where a domain $\mathcal{X} \subset D(F_1) \cap D(F_2)$, the functions

$$P_\xi: x \rightarrow (\nu^{*0\xi}x)^2 + (\tilde{\nu}^{0\xi}x)^2 \quad \text{for all } x \in \mathbb{R}^n, \quad \xi = 1, \dots, k_1, 2k_1 + 1,$$

$$\varphi_\xi: x \rightarrow \operatorname{arctg} \frac{\tilde{\nu}^{0\xi}x}{\nu^{*0\xi}x} \quad \text{for all } x \in \mathcal{X}, \quad \xi = 1, \dots, k_1, 2k_1 + 1,$$

and $h_{q\xi} = \tilde{h}_{q\xi} + \tilde{h}_{q\xi} i$, $h_{q\theta} = \tilde{h}_{q\theta} + \tilde{h}_{q\theta} i$ is a nontrivial solution to the system

$$\sum_{\xi=1}^{2k_1} \left(\lambda_\xi^j h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{j\zeta} h_{q\xi} \right) + \lambda_{2k_1+1}^j h_{0,(2k_1+1)} + \sum_{\theta=1}^{k_2} \left(\lambda_\theta^j h_{0\theta} + \sum_{q=1}^{\varepsilon_\theta} \mu_{q\theta}^{j\zeta} h_{q\theta} \right) = 0, \quad j = 1, \dots, m.$$

Here $\nu^{0\xi} = \nu^{*0\xi} + \tilde{\nu}^{0\xi} i$, $\nu^{0,(k_1+\xi)} = \overline{\nu^{0\xi}}$, $\nu^{0,(2k_1+1)} = \nu^{*0,(2k_1+1)} + \tilde{\nu}^{0,(2k_1+1)} i$ are complex eigenvectors of the matrices A_j corresponding to the eigenvalues $\lambda_\xi^j = \lambda_\xi^{*j} + \tilde{\lambda}_\xi^j i$, $\lambda_{k_1+\xi}^j = \overline{\lambda_\xi^j}$, $\xi = 1, \dots, k_1$, $\lambda_{2k_1+1}^j = \lambda_{2k_1+1}^{*j} + \tilde{\lambda}_{2k_1+1}^j i$, $j = 1, \dots, m$, respectively; $\nu^{0\theta}$ are real common eigenvectors of A_j corresponding to the eigenvalues λ_θ^j , $j = 1, \dots, m$, $\theta = 1, \dots, k_2$, respectively; the functions $v_{q\xi}^\zeta = \tilde{v}_{q\xi}^\zeta + \tilde{v}_{q\xi}^\zeta i$, $v_{q\theta}^\zeta$ is the solution to the system (2.17); the numbers

$$\mu_{q\xi}^{j\zeta} = \mathfrak{p}_j v_{q\xi}^\zeta(x), \quad \mu_{q\xi}^{*j\zeta} = \operatorname{Re} \mu_{q\xi}^{j\zeta}, \quad \tilde{\mu}_{q\xi}^{j\zeta} = \operatorname{Im} \mu_{q\xi}^{j\zeta}, \quad q = 1, \dots, \varepsilon_\xi, \quad \xi = 1, \dots, 2k_1,$$

$$\mu_{q\theta}^{j\zeta} = \mathfrak{p}_j v_{q\theta}^\zeta(x), \quad q = 1, \dots, \varepsilon_\theta, \quad \theta = 1, \dots, k_2, \quad j = 1, \dots, m;$$

the numbers $\varepsilon_\xi, \varepsilon_\theta$ such that $2 \sum_{\xi=1}^{k_1} \varepsilon_\xi + \sum_{\theta=1}^{k_2} \varepsilon_\theta = m - 2k_1 - k_2$ with $2k_1 + 1 + k_2 \leq r$, $\varepsilon_\xi \leq s_\xi - 1$, $\xi = 1, \dots, k_1$, $\varepsilon_\theta \leq s_\theta - 1$, $\theta = 1, \dots, k_2$, where k_1 is a number of complex common eigenvectors (this set hasn't complex conjugate vectors) of the matrices A_j and k_2 is a number of real common eigenvectors of the matrices A_j , $j = 1, \dots, m$.

As an example, the completely solvable system of equations in total differentials

$$\begin{aligned} dx_1 &= (1, -2, 2, 0, 1, 1)xdt_1 + (0, 2, 0, 0, 1, 1)xdt_2 + (3, 0, 0, 0, -1, -1)xdt_3 + (1, -2, 4, 0, 2, 2)xdt_4, \\ dx_2 &= (0, 2, -2, 0, -2, -2)xdt_1 - (1, 3, 0, 0, 1, 1)xdt_2 + (-1, 2, 0, 0, 1, 1)xdt_3 - (2, 1, 4, 0, 4, 4)xdt_4, \\ dx_3 &= (0, 3, -2, 0, -2, -2)xdt_1 - (1, 3, 1, 0, 2, 2)xdt_2 + \\ &\quad + (-2, -1, 2, 0, 1, 1)xdt_3 - (3, -2, 7, 0, 5, 5)xdt_4, \end{aligned} \quad (2.28)$$

$$\begin{aligned} dx_4 &= x((0, -4, 0, 2, -2, 2)dt_1 + (2, 2, 0, 1, 0, 4)dt_2 + (1, 2, -2, 1, -1, -1)dt_3 + (3, -4, 10, 2, 7, 7)dt_4), \\ dx_5 &= (2, -3, 4, 2, 2, 4)xdt_1 + (3, 3, 2, 2, 1, 4)xdt_2 + (2, 1, -1, 0, 0, -1)xdt_3 + (3, -2, 9, 0, 7, 5)xdt_4, \\ dx_6 &= x(-(1, -3, 2, 2, -1, 1)dt_1 - (2, 1, 2, 2, 1, 4)dt_2 + (-1, -1, 1, 0, 1, 2)dt_3 - (-1, -4, 5, 0, 4, 2)dt_4) \end{aligned}$$

has the eigenvalue $\lambda_1^1 = 1 + i$ with elementary divisor $(\lambda^1 - 1 - i)^2$ corresponding to the eigenvector $\nu^{01} = (1, 1 + i, 0, 0, i, i)$ and to the generalized eigenvector $\nu^{11} = (1 + i, 0, 1 + i, 0, i, i)$ and the simple eigenvalue $\lambda_2^1 = 2i$ with common eigenvector $\nu^{02} = (1, 0, 1 + i, 1, i, 1)$.

The real-valued scalar functions

$$\begin{aligned} v_{11}^1: x &\rightarrow \frac{(x_1 + x_2)(x_1 + x_3) + (x_2 + x_5 + x_6)(x_1 + x_3 + x_5 + x_6)}{P_1(x)} \quad \text{for all } x \in \mathcal{X}, \\ \tilde{v}_{11}^1: x &\rightarrow \frac{(x_1 + x_2)(x_1 + x_3 + x_5 + x_6) - (x_1 + x_3)(x_2 + x_5 + x_6)}{P_1(x)} \quad \text{for all } x \in \mathcal{X}, \end{aligned} \quad (2.29)$$

where $P_1: x \rightarrow (x_1 + x_2)^2 + (x_2 + x_5 + x_6)^2$ for all $x \in \mathbb{R}^6$.

The autonomous general integral of the system (2.28) is the functions

$$F_1: x \rightarrow P_1(x)(P_2(x))^2 \exp(-10\varphi_1(x) + 8v_{11}^*(x) + 6\tilde{v}_{11}^1(x)) \quad \text{for all } x \in \mathcal{X} \quad (2.30)$$

and

$$F_2: x \rightarrow (P_1(x))^3 \exp(-10\varphi_1(x) - 4\varphi_2(x) + 12v_{11}^*(x) + 14\tilde{v}_{11}^1(x)) \quad \text{for all } x \in \mathcal{X}, \quad (2.31)$$

where a domain $\mathcal{X} \subset \{x: x_1 + x_2 \neq 0, x_1 + x_3 + x_4 + x_6 \neq 0\}$, the scalar functions

$$P_2: x \rightarrow (x_1 + x_3 + x_4 + x_6)^2 + (x_3 + x_5)^2 \quad \text{for all } x \in \mathbb{R}^6,$$

$$\varphi_1: x \rightarrow \operatorname{arctg} \frac{x_2 + x_5 + x_6}{x_1 + x_2}, \quad \varphi_2: x \rightarrow \operatorname{arctg} \frac{x_3 + x_5}{x_1 + x_3 + x_4 + x_6} \quad \text{for all } x \in \mathcal{X}.$$

Case 2B. A function $v_{l\gamma}^\zeta$, $\gamma \in \{1, \dots, k_1\}$, $l \in \{1, \dots, \varepsilon_\gamma\}$ hasn't the complex conjugate function. Autonomous first integrals of the completely solvable system (2.1) are the functions

$$F_1: x \rightarrow \prod_{\xi=1}^{k_1} (P_\xi(x))^{h_{0\xi}^* + h_{0,(k_1+\xi)}^*} \exp\left(-2(\tilde{h}_{0\xi} - \tilde{h}_{0,(k_1+\xi)}) \varphi_\xi(x) + \right.$$

$$+ 2 \sum_{q=1}^{\varepsilon_\xi} (1 - \delta_{ql} \delta_{\xi\gamma}) \left((h_{q\xi}^* + h_{q,(k_1+\xi)}^*) v_{q\xi}^\zeta(x) + (\tilde{h}_{q,(k_1+\xi)} - \tilde{h}_{q\xi}) \tilde{v}_{q\xi}^\zeta(x) \right) +$$

$$+ 2 \left(h_{l\gamma}^* v_{l\gamma}^\zeta(x) - \tilde{h}_{l\gamma} \tilde{v}_{l\gamma}^\zeta(x) \right) \prod_{\theta=1}^{k_2} (\nu^{0\theta} x)^{2h_{0\theta}^*} \exp\left(2 \sum_{q=1}^{\varepsilon_\theta} h_{q\theta}^* v_{q\theta}^\zeta(x)\right) \quad \text{for all } x \in \mathcal{X},$$

and

$$F_2: x \rightarrow \prod_{\xi=1}^{k_1} (P_\xi(x))^{\tilde{h}_{0\xi} + \tilde{h}_{0,(k_1+\xi)}} \exp\left(2(h_{0\xi}^* - h_{0,(k_1+\xi)}^*) \varphi_\xi(x) + \right.$$

$$+ 2 \sum_{q=1}^{\varepsilon_\xi} (1 - \delta_{ql} \delta_{\xi\gamma}) \left((\tilde{h}_{q\xi} + \tilde{h}_{q,(k_1+\xi)}) v_{q\xi}^\zeta(x) + (h_{q\xi}^* - h_{q,(k_1+\xi)}^*) \tilde{v}_{q\xi}^\zeta(x) \right) +$$

$$+ 2(h_{l\gamma}^* \tilde{v}_{l\gamma}^\zeta(x) + \tilde{h}_{l\gamma} v_{l\gamma}^\zeta(x)) \prod_{\theta=1}^{k_2} (\nu^{0\theta} x)^{2\tilde{h}_{0\theta}} \exp\left(2 \sum_{q=1}^{\varepsilon_\theta} \tilde{h}_{q\theta} v_{q\theta}^\zeta(x)\right) \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset D(F_1) \cap D(F_2)$, δ is the Kronecker delta, the functions

$$P_\xi: x \rightarrow (\nu^{0\xi} x)^2 + (\tilde{\nu}^{0\xi} x)^2 \quad \text{for all } x \in \mathbb{R}^n, \quad \varphi_\xi: x \rightarrow \operatorname{arctg} \frac{\tilde{\nu}^{0\xi} x}{\nu^{0\xi} x} \quad \text{for all } x \in \mathcal{X}, \quad \xi = 1, \dots, k_1,$$

and $h_{q\xi} = h_{q\xi}^* + \tilde{h}_{q\xi} i$, $h_{q\theta} = h_{q\theta}^* + \tilde{h}_{q\theta} i$ is a nontrivial solution to the linear system

$$\sum_{\xi=1}^{2k_1} \left(\lambda_\xi^j h_{0\xi} + \sum_{q=1}^{\varepsilon_\xi} \mu_{q\xi}^{j\zeta} h_{q\xi} \right) - \mu_{l,(k_1+\gamma)}^{j\zeta} h_{l,(k_1+\gamma)} + \sum_{\theta=1}^{k_2} \left(\lambda_\theta^j h_{0\theta} + \sum_{q=1}^{\varepsilon_\theta} \mu_{q\theta}^{j\zeta} h_{q\theta} \right) = 0, \quad j = 1, \dots, m.$$

Here $\nu^{0\xi} = \nu^{0\xi} + \tilde{\nu}^{0\xi} i$, $\nu^{0,(k_1+\xi)} = \overline{\nu^{0\xi}}$ are complex common eigenvectors of the matrices A_j corresponding to the eigenvalues $\lambda_\xi^j = \lambda_\xi^j + \tilde{\lambda}_\xi^j i$, $\lambda_{k_1+\xi}^j = \overline{\lambda_\xi^j}$, $j = 1, \dots, m$, $\xi = 1, \dots, k_1$, respectively; $\nu^{0\theta}$ are real common eigenvectors of the matrices A_j corresponding to the eigenvalues λ_θ^j , $j = 1, \dots, m$, $\theta = 1, \dots, k_2$, respectively; the functions $v_{q\xi}^\zeta = v_{q\xi}^* + \tilde{v}_{q\xi} i$, $v_{q\theta}^\zeta$

is the solution to the system (2.17); the numbers

$$\mu_{q\xi}^{j\xi} = \mathfrak{p}_j v_{q\xi}^\zeta(x), \quad \mu_{q\xi}^{*j\xi} = \operatorname{Re} \mu_{q\xi}^{j\xi}, \quad \tilde{\mu}_{q\xi}^{j\xi} = \operatorname{Im} \mu_{q\xi}^{j\xi}, \quad q = 1, \dots, \varepsilon_\xi, \quad \xi = 1, \dots, 2k_1,$$

$$\mu_{q\theta}^{j\zeta} = \mathfrak{p}_j v_{q\theta}^\zeta(x), \quad q = 1, \dots, \varepsilon_\theta, \quad \theta = 1, \dots, k_2, \quad j = 1, \dots, m;$$

the numbers $\varepsilon_\xi, \varepsilon_\theta$ such that $2 \sum_{\xi=1}^{k_1} \varepsilon_\xi + \sum_{\theta=1}^{k_2} \varepsilon_\theta = m - 2k_1 - k_2 + 2$ with $2k_1 + k_2 \leq r$, $\varepsilon_\xi \leq s_\xi - 1$, $\xi = 1, \dots, k_1$, $\varepsilon_\theta \leq s_\theta - 1$, $\theta = 1, \dots, k_2$, where k_1 is a number of complex common eigenvectors (this set hasn't complex conjugate vectors) of the matrices A_j and k_2 is a number of real common eigenvectors of the matrices A_j , $j = 1, \dots, m$.

In particular, the completely solvable system of total differential equations

$$\begin{aligned} dx_1 &= (3, -4, 4, 1, 0, 2)x dt_1 + (0, -4, 2, 1, -1, 1)x dt_2, \\ dx_2 &= (-1, 3, -3, 0, -2, -3)x dt_1 + (1, 3, 0, 0, 1, -1)x dt_2, \\ dx_3 &= (-3, 5, -5, -1, -2, -4)x dt_1 + (0, 6, -2, -1, 2, -1)x dt_2, \\ dx_4 &= (3, -6, 4, 4, -1, 5)x dt_1 + (2, -6, 2, 3, -4, 2)x dt_2, \\ dx_5 &= (5, -5, 8, 3, 3, 6)x dt_1 + (1, -6, 3, 2, -2, 2)x dt_2, \\ dx_6 &= (-2, 5, -4, -3, 1, -2)x dt_1 + (-2, 4, -3, -3, 2, -2)x dt_2 \end{aligned} \quad (2.32)$$

has the eigenvalue $\lambda_1^1 = 1 + 2i$ with elementary divisor $(\lambda^1 - 1 - 2i)^3$ corresponding to the eigenvector $\nu^{01} = (1, 0, 1 + i, 1, i, 1)$ and to the generalized eigenvectors $\nu^{11} = (1, 1 + i, 0, 0, i, i)$, $\nu^{21} = (2 + 2i, 0, 2 + 2i, 0, 2i, 2i)$. Autonomous first integrals of the system (2.32) are the functions

$$F_1: x \rightarrow P(x) \exp(-\varphi(x) - \tilde{v}_{11}^1(x)) \quad \text{for all } x \in \mathcal{X}, \quad (2.33)$$

$$F_2: x \rightarrow P(x) \exp(-2\varphi(x) + 2\tilde{v}_{11}^{*1}(x)) \quad \text{for all } x \in \mathcal{X}, \quad (2.34)$$

$$F_3: x \rightarrow P^2(x) \exp(-2\varphi(x) - \tilde{v}_{21}^1(x)) \quad \text{for all } x \in \mathcal{X}, \quad (2.35)$$

and

$$F_4: x \rightarrow \tilde{v}_{21}^{*1}(x) \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 + x_3 + x_4 + x_6 \neq 0\}, \quad (2.36)$$

where the functions $\tilde{v}_{11}^1, \tilde{v}_{11}^{*1}, \tilde{v}_{21}^1, \tilde{v}_{21}^{*1}, P$, and φ are given by (2.22), (2.23), and (2.27).

Remark 2.1 Consider the completely solvable system of equations in total differentials

$$\begin{aligned} dx_1 &= (2x_1 + x_3) dt_1 + (2x_1 + 3x_2 + 3x_3) dt_2, \quad dx_2 = (x_1 + x_2 + x_3) dt_1 + 2x_2 dt_2, \\ dx_3 &= -x_1 dt_1 - (3x_2 + x_3) dt_2, \quad dx_4 = x_4 dt_1 + (x_2 + x_3 - x_4) dt_2. \end{aligned} \quad (2.37)$$

The matrices A_1 and A_2 of (2.37) have the elementary divisors $(\lambda^1 - 1)^2, \lambda^1 - 1, \lambda^1 - 1$, and $\lambda^2 - 2, \lambda^2 - 2, (\lambda^2 + 1)^2$, respectively. Using the common eigenvector $\nu^{01} = (1, 0, 1, 0)$, generalized eigenvector $\nu^{11} = (0, 1, 0, 0)$ of A_1 and the common eigenvector $\nu^{03} = (0, 1, 1, 0)$, generalized eigenvector $\nu^{13} = (0, 0, 0, 1)$ of the matrix A_2 , we can build the functions

$$v_{11}^1: x \rightarrow \frac{x_2}{x_1 + x_3}, \quad v_{13}^2: x \rightarrow \frac{x_4}{x_2 + x_3} \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 + x_3 \neq 0, x_2 + x_3 \neq 0\}.$$

The system (2.37) has the autonomous first integrals on a domain $\mathcal{X}$:

$$F_1: (t, x) \rightarrow (x_1 + x_3)(x_2 + x_3)^2 \exp(-3v_{11}^1(x)), \quad F_2: (t, x) \rightarrow \frac{x_2 + x_3}{x_1 + x_3} \exp(3v_{13}^2(x)).$$

2.1.3. Nonautonomous first integrals.

Theorem 2.5. Suppose ν is a real common eigenvector of the matrices A_j corresponding to the eigenvalues λ^j , $j = 1, \dots, m$, respectively. Then the scalar function

$$F: (t, x) \rightarrow (\nu x) \exp\left(-\sum_{j=1}^m \lambda^j t_j\right) \quad \text{for all } (t, x) \in \mathbb{R}^{n+m}$$

is a first integral of the linear system of equations in total differentials (2.1).

Consider the system (2.3). Using the eigenvalues $\lambda_1^1 = -2$, $\lambda_1^2 = 1$ corresponding to the common eigenvector $\nu^1 = (0, -1, 1, 1)$ and the eigenvalues $\lambda_2^1 = 0$, $\lambda_2^2 = -1$ corresponding to the eigenvector $\nu^2 = (1, 0, 0, 0)$, we can build the first integrals of the system (2.3):

$$F_1: (t, x) \rightarrow (-x_2 + x_3 + x_4) \exp(2t_1 - t_2), \quad F_2: (t, x) \rightarrow x_1 \exp t_2 \quad \text{for all } (t, x) \in \mathbb{R}^6.$$

The functions F_1 , F_2 , (2.4), and (2.5) are the general integral of the system (2.3).

Corollary 2.2. Let $\nu = \nu^* + \tilde{\nu}i$ ($\nu^* = \operatorname{Re} \nu$, $\tilde{\nu} = \operatorname{Im} \nu$) be a common complex eigenvector of the matrices A_j corresponding to the eigenvalues $\lambda^j = \lambda^{*j} + \tilde{\lambda}^j i$ ($\lambda^{*j} = \operatorname{Re} \lambda^j$, $\tilde{\lambda}^j = \operatorname{Im} \lambda^j$), $j = 1, \dots, m$, respectively. Then the system (2.1) has the first integrals

$$F_1: (t, x) \rightarrow ((\nu^* x)^2 + (\tilde{\nu} x)^2) \exp\left(-2 \sum_{j=1}^m \lambda^{*j} t_j\right) \quad \text{for all } (t, x) \in \mathbb{R}^{n+m}$$

and

$$F_2: (t, x) \rightarrow \operatorname{arctg} \frac{\tilde{\nu} x}{\nu^* x} - \sum_{j=1}^m \tilde{\lambda}^j t_j \quad \text{for all } (t, x) \in \mathcal{D}, \quad \mathcal{D} \subset \mathbb{R}^{n+m}.$$

For example, the completely solvable system (2.8) has the eigenvector $\nu^1 = (1, i, 0)$ corresponding to the eigenvalues $\lambda_1^1 = 1$, $\lambda_1^2 = -i$ and the first integrals (by Corollary 2.2)

$$F_1: (t, x) \rightarrow (x_1^2 + x_2^2) \exp(-2t_1) \quad \text{for all } (t, x) \in \mathbb{R}^5,$$

$$F_2: (t, x) \rightarrow \operatorname{arctg} \frac{x_2}{x_1} + t_2 \quad \text{for all } (t, x) \in \mathbb{R}^2 \times \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 \neq 0, x_3 \neq 0\}.$$

The functionally independent first integrals (2.9), F_1 , and F_2 are the general integral on a domain $\mathbb{R}^2 \times \mathcal{X}$ for the system of equations in total differentials (2.8).

The completely solvable system (2.12) has the eigenvector $\nu^1 = (0, -i, 0, 1)$ corresponding to the eigenvalues $\lambda_1^1 = -1$ and $\lambda_1^2 = -i$. The first integrals (2.13) and (by Corollary 2.2)

$$F_1: (t, x) \rightarrow (x_2^2 + x_4^2) \exp(2t_1) \quad \text{for all } (t, x) \in \mathbb{R}^6,$$

$$F_2: (t, x) \rightarrow \operatorname{arctg} \frac{x_2}{x_4} - t_2 \quad \text{for all } (t, x) \in \mathbb{R}^2 \times \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 x_4 - x_2 x_3 \neq 0, x_4 \neq 0\},$$

are the general integral on a domain $\mathbb{R}^2 \times \mathcal{X}$ of the linear system (2.12).

Theorem 2.6. Let ν^0 be a real common eigenvector of the matrices A_j corresponding to the eigenvalues λ^j , $j = 1, \dots, m$, respectively. Let ν^θ , $\theta = 1, \dots, s-1$ be real generalized eigenvectors of the matrix A_ζ corresponding to the eigenvalue λ^ζ with elementary divisor of multiplicity $s \geq 2$. Then the completely solvable system (2.1) has the first integrals

$$F_q: (t, x) \rightarrow v_q^\zeta(x) - \sum_{j=1}^m \mu_q^{j\zeta} t_j \quad \text{for all } (t, x) \in \mathbb{R}^m \times \mathcal{X}, \quad q = 1, \dots, s-1,$$

where the set of functions $v_q^\zeta: \mathcal{X} \rightarrow \mathbb{R}$ is the solution to the system (2.17), the numbers $\mu_q^{j\zeta} = \mathfrak{p}_j v_q^\zeta(x)$, $q = 1, \dots, s-1$, $j = 1, \dots, m$, and a domain $\mathcal{X} \subset \{x: \nu^0 x \neq 0\}$.

The completely solvable system (2.18) has $\mu_1^{11} = 1$, $\mu_3^{11} = 0$, $\mu_1^{21} = -1$, $\mu_3^{21} = 6$, and the first integrals (by Theorem 2.6)

$$F_1: (t, x) \rightarrow v_{11}^1(x) - t_1 + t_2 \quad \text{for all } (t, x) \in \mathbb{R}^2 \times \mathcal{X},$$

$$F_2: (t, x) \rightarrow v_{31}^1(x) - 6t_2 \quad \text{for all } (t, x) \in \mathbb{R}^2 \times \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 - x_2 + x_3 \neq 0\},$$

where the functions v_{11}^1, v_{31}^1 are given by (2.19). The functionally independent first integrals (2.20), F_1 , and F_2 are the general integral on a domain $\mathbb{R}^2 \times \mathcal{X}$ of the system (2.18).

Corollary 2.3. *Let ν^0 be a common eigenvector of the matrices A_j corresponding to the eigenvalues λ^j , $j = 1, \dots, m$. Suppose ν^θ , $\theta = 1, \dots, s-1$ are generalized eigenvectors of A_ζ corresponding to the complex eigenvalue λ^ζ ($\text{Im } \lambda^\zeta \neq 0$) with elementary divisor of multiplicity $s \geq 2$. Then the completely solvable system (2.1) has the first integrals*

$$F_{1q}: (t, x) \rightarrow v_q^\zeta(x) - \sum_{j=1}^m \mu_q^{j\zeta} t_j \quad \text{for all } (t, x) \in \mathbb{R}^m \times \mathcal{X}, \quad q = 1, \dots, s-1,$$

and

$$F_{2q}: (t, x) \rightarrow \tilde{v}_q^\zeta(x) - \sum_{j=1}^m \tilde{\mu}_q^{j\zeta} t_j \quad \text{for all } (t, x) \in \mathbb{R}^m \times \mathcal{X}, \quad q = 1, \dots, s-1, \quad \mathcal{X} \subset \mathbb{R}^n,$$

where the set of functions $v_q^\zeta: x \rightarrow v_q^\zeta(x) + \tilde{v}_q^\zeta(x)i$ for all $x \in \mathcal{X}$ is the solution to the system (2.17), the numbers $\mu_q^{j\zeta} = \mathbf{p}_j v_q^\zeta(x)$, $\tilde{\mu}_q^{j\zeta} = \mathbf{p}_j \tilde{v}_q^\zeta(x)$, $j = 1, \dots, m$, $q = 1, \dots, s-1$.

The completely solvable system (2.21) has the first integrals (by Corollary 2.3)

$$F_{11}: (t, x) \rightarrow v_{11}^1(x) - t_1 - t_2, \quad F_{21}: (t, x) \rightarrow \tilde{v}_{11}^1(x) + t_2 - t_3 \quad \text{for all } (t, x) \in \mathbb{R}^3 \times \mathcal{X},$$

and

$$F_{12}: (t, x) \rightarrow v_{21}^1(x) - 2t_3 \quad \text{for all } (t, x) \in \mathbb{R}^3 \times \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 + x_3 + x_4 + x_6 \neq 0\},$$

where $v_{11}^1, \tilde{v}_{11}^1$, and v_{21}^1 are given by (2.22). The general integral for the linear system (2.21) is the functionally independent first integrals (2.24), (2.25), (2.26), F_{11} , F_{21} , and F_{12} .

For example, the system (2.28) has the eigenvector $\nu^{01} = (1, 1+i, 0, 0, i, i)$ corresponding to the eigenvalues $\lambda_1^1 = 1+i$, $\lambda_1^2 = -1$, $\lambda_1^3 = 2$, $\lambda_1^4 = -1+2i$ and the first integrals

$$F_1: (t, x) \rightarrow ((x_1 + x_2)^2 + (x_2 + x_5 + x_6)^2) \exp(-2t_1 + 2t_2 - 4t_3 + 2t_4) \quad (\text{by Corollary 2.2}),$$

$$F_2: (t, x) \rightarrow \arctg \frac{x_2 + x_5 + x_6}{x_1 + x_2} - t_1 - 2t_4 \quad \text{for all } (t, x) \in \mathbb{R}^4 \times \mathcal{X},$$

where $\mathcal{X} \subset \{x: x_1 + x_2 \neq 0, x_1 + x_3 + x_4 + x_6 \neq 0\} \subset \mathbb{R}^6$. Using Corollary 2.3, we get

$$F_3: (t, x) \rightarrow v_{11}^1(x) - t_1 + t_3 - t_4 \quad \text{for all } (t, x) \in \mathbb{R}^4 \times \mathcal{X}$$

and

$$F_4: (t, x) \rightarrow \tilde{v}_{11}^1(x) - t_2 - t_4 \quad \text{for all } (t, x) \in \mathbb{R}^4 \times \mathcal{X},$$

where v_{11}^1 and $\tilde{v}_{11}^1$ are given by (2.29). The first integrals (2.30), (2.31), $F_1, \dots, F_4$ are the general integral on a domain $\mathbb{R}^4 \times \mathcal{X}$ for the completely solvable linear system (2.28).

Consider the completely solvable linear system of equations in total differentials (2.32). According to Corollary 2.3, we can construct the first integrals

$$F_{11}: (t, x) \rightarrow v_{11}^1(x) - t_1 - t_2, \quad F_{21}: (t, x) \rightarrow \tilde{v}_{11}^1(x) + t_2 \quad \text{for all } (t, x) \in \mathbb{R}^2 \times \mathcal{X},$$

where the functions v_{11}^1 and $\tilde{v}_{11}^1$ are given by (2.22), a domain $\mathcal{X} \subset \{x: x_1 + x_3 + x_4 + x_6 \neq 0\}$.

The functionally independent first integrals (2.33), (2.34), (2.35), (2.36), F_{11} , and F_{21} are the general integral on a domain $\mathbb{R}^2 \times \mathcal{X}$ for the completely solvable system (2.32).

Remark 2.2. The system of equations in total differentials

$$dx_1 = -(x_1 + x_2 + 3x_3) dt_1 - 3(x_1 + 2x_3) dt_2, \quad (2.38)$$

$$dx_2 = x_2 dt_1 - 2x_2 dt_2, \quad dx_3 = (x_2 + 2x_3) dt_1 + 3x_3 dt_2$$

is not completely solvable (this system has the defect [5, p. 54] of order 1):

$$\begin{aligned} [\mathbf{p}_1(x), \mathbf{p}_2(x)] &= [-(x_1 + x_2 + 3x_3) \partial_{x_1} + x_2 \partial_{x_2} + (x_2 + 2x_3) \partial_{x_3}, \\ &- 3(x_1 + 2x_3) \partial_{x_1} - 2x_2 \partial_{x_2} + 3x_3 \partial_{x_3}] = -5x_2 \partial_{x_1} + 5x_2 \partial_{x_3} \equiv \mathbf{p}_3(x) \quad \text{for all } x \in \mathbb{R}^3, \\ [\mathbf{p}_1(x), \mathbf{p}_3(x)] &= -\mathbf{p}_3(x), \quad [\mathbf{p}_2(x), \mathbf{p}_3(x)] = -5\mathbf{p}_3(x) \quad \text{for all } x \in \mathbb{R}^3. \end{aligned}$$

The system (2.38) has two common eigenvectors $\nu^1 = (1, 0, 1)$ and $\nu^2 = (0, 1, 0)$ corresponding to the eigenvalues $\lambda_1^1 = -1$, $\lambda_1^2 = -3$, and $\lambda_2^1 = 1$, $\lambda_2^2 = -2$. The first integrals

$$F_1: (t, x) \rightarrow (x_1 + x_3) \exp(t_1 + 3t_2), \quad F_2: (t, x) \rightarrow x_2 \exp(2t_2 - t_1) \quad \text{for all } (t, x) \in \mathbb{R}^5$$

are a general integral of the linear system of equations in total differentials (2.38).

2.2. Linear real nonhomogeneous systems in total differentials

Consider now a nonhomogeneous system of equations in total differentials

$$dx = \sum_{j=1}^m (B_j x + f_j(t)) dt_j \quad (2.39)$$

corresponding to the linear homogeneous system (2.1), where the matrices $B_1, \dots, B_m$ are transpose of $A_1, \dots, A_m$, respectively, and the vector functions

$$f_j: t \rightarrow \text{colon}(f_{j1}(t), \dots, f_{jn}(t)) \quad \text{for all } t \in \mathcal{T}, \quad j = 1, \dots, m$$

are continuously differentiable on a domain $\mathcal{T} \subset \mathbb{R}^m$.

The Frobenius conditions for the total solvability [4, p. 44] of system (2.39) are (2.2) and

$$\partial_{t_\zeta} f_j(t) - B_\zeta f_j(t) = \partial_{t_j} f_\zeta(t) - B_j f_\zeta(t) \quad \text{for all } t \in \mathcal{T}, \quad j, \zeta = 1, \dots, m. \quad (2.40)$$

Theorem 2.7. *Let ν be a real common eigenvector of the matrices A_j corresponding to the eigenvalues λ^j , $j = 1, \dots, m$, respectively. Then the completely solvable system of equations in total differentials (2.39) has the first integral*

$$F: (t, x) \rightarrow \nu x \cdot \varphi(t) - \int \sum_{j=1}^m \nu f_j(t) \cdot \varphi(t) dt_j \quad \text{for all } (t, x) \in \tilde{\mathcal{T}} \times \mathbb{R}^n, \quad (2.41)$$

where $\tilde{\mathcal{T}}$ is a simply connected domain from $\mathcal{T}$, the function

$$\varphi: t \rightarrow \exp\left(-\sum_{j=1}^m \lambda^j t_j\right) \quad \text{for all } t \in \mathcal{T}.$$

Proof. Let us introduce the 1-form

$$\omega: t \rightarrow \sum_{j=1}^m \nu f_j(t) \cdot \varphi(t) dt_j \quad \text{for all } t \in \mathcal{T}, \quad \mathcal{T} \subset \mathbb{R}^m.$$

The external differential has the form

$$d\omega(t) = \varphi(t) \sum_{j=1}^m \sum_{\zeta=1}^m (\partial_{t_\zeta} \nu f_j(t) - \lambda^\zeta \cdot \nu f_j(t)) dt_\zeta \wedge dt_j \quad \text{for all } t \in \mathcal{T}.$$

Using (2.40) and

$$dt_j \wedge dt_j = 0, \quad dt_\zeta \wedge dt_j = -dt_j \wedge dt_\zeta, \quad \zeta, j = 1, \dots, m,$$

we get the differential 1-form ω is closed on $\mathcal{T}$. By the Poincaré theorem (see [5, p. 14]), it follows that ω is exact in a simply connected domain $\tilde{\mathcal{T}} \subset \mathcal{T}$.

Taking into account Lemma 2.1 and (2.15), we obtain

$$\mathfrak{P}_j F(t, x) = -\lambda^j \nu x \cdot \varphi(t) - \nu f_j(t) \varphi(t) + \mathfrak{p}_j \nu x \cdot \varphi(t) + \nu f_j(t) \varphi(t) = 0$$

$$\text{for all } (t, x) \in \tilde{\mathcal{T}} \times \mathbb{R}^n, \quad j = 1, \dots, m,$$

where the linear differential operators

$$\mathfrak{P}_j(t, x) = \partial_{t_j} + \mathfrak{p}_j(x) + f_j(t) \partial_x \quad \text{for all } (t, x) \in \mathcal{T} \times \mathbb{R}^n, \quad j = 1, \dots, m.$$

Therefore the function (2.41) is a first integral of the completely solvable system (2.39). ■

Corollary 2.4. *Let $\nu = \tilde{\nu}^* + \tilde{\nu} i$ ($\tilde{\nu}^* = \operatorname{Re} \nu$, $\tilde{\nu} = \operatorname{Im} \nu$) be a common complex eigenvector of the matrices A_j corresponding to the eigenvalues $\lambda^j = \tilde{\lambda}^j + \tilde{\lambda}^j i$ ($\tilde{\lambda}^j = \operatorname{Re} \lambda^j$, $\tilde{\lambda}^j = \operatorname{Im} \lambda^j$), $j = 1, \dots, m$, respectively. Then the completely solvable system (2.39) has the first integrals*

$$F_\tau: (t, x) \rightarrow \alpha_\tau(t, x) - \int \sum_{j=1}^m \alpha_\tau(t, f_j(t)) dt_j \quad \text{for all } (t, x) \in \tilde{\mathcal{T}} \times \mathbb{R}^n, \quad \tau = 1, 2,$$

where the scalar functions of the vector arguments

$$\begin{aligned} \alpha_1(t, x) &= \left(\tilde{\nu}^* x \cdot \cos \sum_{j=1}^m \tilde{\lambda}^j t_j + \tilde{\nu} x \cdot \sin \sum_{j=1}^m \tilde{\lambda}^j t_j \right) \cdot \exp \left(- \sum_{j=1}^m \tilde{\lambda}^j t_j \right) \quad \text{for all } (t, x) \in \mathcal{T} \times \mathbb{R}^n, \\ \alpha_2(t, x) &= \left(\tilde{\nu} x \cdot \cos \sum_{j=1}^m \tilde{\lambda}^j t_j - \tilde{\nu}^* x \cdot \sin \sum_{j=1}^m \tilde{\lambda}^j t_j \right) \cdot \exp \left(- \sum_{j=1}^m \tilde{\lambda}^j t_j \right) \quad \text{for all } (t, x) \in \mathcal{T} \times \mathbb{R}^n. \end{aligned}$$

For example, the completely solvable system of total differential equations

$$\begin{aligned} dx_1 &= x_1 dt_1 + (x_2 + 1) dt_2, & dx_2 &= (x_2 + 1) dt_1 + (-x_1 + e^{t_1 + at_2}) dt_2, \\ dx_3 &= (x_3 + t_2 - t_1) dt_1 + (-x_3 + t_1 - t_2) dt_2 \end{aligned}$$

corresponding to the linear homogeneous system (2.8). This system has the first integrals

$$F_1: (t, x) \rightarrow (x_1 \cos t_2 - x_2 \sin t_2 - \sin t_2) e^{-t_1} + \frac{a \sin t_2 - \cos t_2}{a^2 + 1} e^{at_2} \quad (\text{by Corollary 2.4}),$$

$$F_2: (t, x) \rightarrow (x_1 \sin t_2 + x_2 \cos t_2 + \cos t_2) e^{-t_1} - \frac{a \cos t_2 + \sin t_2}{a^2 + 1} e^{at_2} \quad (\text{by Corollary 2.4}),$$

$$F_3: (t, x) \rightarrow (x_3 + t_2 - t_1 - 1) e^{t_2 - t_1} \quad \text{for all } (t, x) \in \mathbb{R}^5 \quad (\text{by Theorem 2.7}),$$

where a is some real number.

Theorem 2.8. *Let the assumptions of Lemma 2.2 hold. Then the completely solvable system of equations in total differentials (2.39) has the first integrals*

$$\begin{aligned} F_\theta: (t, x) &\rightarrow \nu^{\theta l} x \cdot \varphi(t) - \sum_{\tau=1}^{\theta} K_{\tau-1}^\theta(t) \cdot F_{\tau-1}(t, x) - C_\theta(t) \\ &\text{for all } (t, x) \in \tilde{\mathcal{T}} \times \mathcal{X}, \quad \theta = 0, \dots, s_l - 1, \quad \mathcal{X} \subset \{x: \nu^0 x \neq 0\}, \end{aligned} \tag{2.42}$$

where the functions

$$\varphi: t \rightarrow \exp\left(-\sum_{j=1}^m \mu_{0l}^{j\zeta} t_j\right) \quad \text{for all } t \in \tilde{T}, \quad \tilde{T} \subset \mathcal{T},$$

$$K_{\tau-1}^\theta: t \rightarrow \int \sum_{j=1}^m \left(\binom{\theta}{\tau-1} \mu_{\theta-\tau+1,l}^{j\zeta} + \sum_{\delta=1}^{\theta-\tau} \binom{\theta}{\delta} \mu_{\delta l}^{j\zeta} \cdot K_{\tau-1}^{\theta-\delta}(t) \right) dt_j, \quad \tau = 1, \dots, \theta, \quad \theta = 1, \dots, s_l - 1,$$

$$C_\theta: t \rightarrow \int \sum_{j=1}^m \left(\nu^{\theta l} f_j(t) \cdot \varphi(t) + \sum_{\tau=1}^{\theta} \binom{\theta}{\tau} \mu_{\tau l}^{j\zeta} \cdot C_{\theta-\tau}(t) \right) dt_j \quad \text{for all } t \in \tilde{T}, \quad \theta = 0, \dots, s_l - 1,$$

the numbers $\mu_{0l}^{j\zeta} = \lambda_l^j$, $\mu_{\theta l}^{j\zeta} = \mathfrak{p}_j v_{\theta l}^\zeta(x)$, $\theta = 1, \dots, s_l - 1$, $j = 1, \dots, m$.

Proof. The proof is by induction on θ .

The case $\theta = 0$ was considered in Theorem 2.7.

Suppose $\theta = 1$. Using Lemma 2.3 with condition (2.16), we obtain

$$\begin{aligned} \mathfrak{P}_j F_1(t, x) &= \mathfrak{P}_j \left(\nu^{1l} x \cdot \varphi(t) - \int \sum_{\xi=1}^m \mu_{1l}^{\xi\zeta} dt_\xi \cdot F_0(t, x) - \int \sum_{\xi=1}^m (\nu^{1l} f_\xi(t) \cdot \varphi(t) + \mu_{1l}^{\xi\zeta} C_0(t)) dt_\xi \right) = \\ &= \mu_{1l}^{j\zeta} (\nu^{0l} x \cdot \varphi(t) - C_0(t) - F_0(t, x)) = 0 \quad \text{for all } (t, x) \in \tilde{T} \times \mathcal{X}, \quad j = 1, \dots, m. \end{aligned}$$

Therefore $F_1: \tilde{T} \times \mathcal{X} \rightarrow \mathbb{R}$ is a first integral of the completely solvable system (2.39).

Suppose that the assertion of the theorem is valid for $\theta = \varepsilon - 1$, i.e., the scalar functions $F_\theta: \tilde{T} \times \mathcal{X} \rightarrow \mathbb{R}$, $\theta = 1, \dots, \varepsilon - 1$ are first integrals of the system (2.39). Then

$$\begin{aligned} \mathfrak{P}_j F_\varepsilon(t, x) &= \mathfrak{P}_j \left(\nu^{\varepsilon l} x \cdot \varphi(t) - \sum_{\tau=1}^{\varepsilon} K_{\tau-1}^\varepsilon(t) \cdot F_{\tau-1}(t, x) - C_\varepsilon(t) \right) = \\ &= -\mu_{0l}^{j\zeta} \nu^{\varepsilon l} x \cdot \varphi(t) + \sum_{\tau=0}^{\varepsilon} \binom{\varepsilon}{\tau} \mu_{\tau l}^{j\zeta} \nu^{\varepsilon-\tau, l} x \cdot \varphi(t) + \nu^{\varepsilon l} f_j(t) \cdot \varphi(t) - \\ &- \sum_{\tau=1}^{\varepsilon} \left(\binom{\varepsilon}{\tau-1} \mu_{\varepsilon-\tau+1,l}^{j\zeta} + \sum_{\delta=1}^{\varepsilon-\tau} \binom{\varepsilon}{\delta} \mu_{\delta l}^{j\zeta} K_{\tau-1}^{\varepsilon-\delta}(t) \right) F_{\tau-1}(t, x) - \left(\nu^{\varepsilon l} f_j(t) \cdot \varphi(t) + \sum_{\tau=1}^{\varepsilon} \binom{\varepsilon}{\tau} \mu_{\tau l}^{j\zeta} C_{\varepsilon-\tau}(t) \right) = \\ &= \sum_{\tau=1}^{\varepsilon} \binom{\varepsilon}{\tau} \mu_{\tau l}^{j\zeta} \left(\left(\nu^{\varepsilon-\tau, l} x \cdot \varphi(t) - \sum_{\delta=1}^{\varepsilon-\tau} K_{\delta-1}^{\varepsilon-\tau}(t) \cdot F_{\delta-1}(t, x) - C_{\varepsilon-\tau}(t) \right) - F_{\varepsilon-\tau}(t, x) \right) = 0 \\ &\quad \text{for all } (t, x) \in \tilde{T} \times \mathcal{X}, \quad j = 1, \dots, m. \end{aligned}$$

Therefore $F_\varepsilon: \tilde{T} \times \mathcal{X} \rightarrow \mathbb{R}$ is a first integral of the linear system (2.39).

Thus the functions (2.42) are first integrals of the completely solvable system (2.39). ■

Remark 2.3. In complex case from Theorem 2.8, we get the following real-valued first integrals of the completely solvable system of equations in total differentials (2.39):

$$F_\theta^1: (t, x) \rightarrow \operatorname{Re} F_\theta(t, x), \quad F_\theta^2: (t, x) \rightarrow \operatorname{Im} F_\theta(t, x) \quad \text{for all } (t, x) \in \tilde{T} \times \mathcal{X}, \quad \theta = 0, \dots, s_l - 1.$$

Moreover, we have

$$F_\theta^1: (t, x) \rightarrow \alpha_\theta(t, x) - \sum_{\tau=1}^{\theta} \left(\operatorname{Re} K_{\tau-1}^\theta(t) \cdot F_{\tau-1}^1(t, x) - \operatorname{Im} K_{\tau-1}^\theta(t) \cdot F_{\tau-1}^2(t, x) \right) - \operatorname{Re} C_\theta(t),$$

$$F_\theta^2: (t, x) \rightarrow \beta_\theta(t, x) - \sum_{\tau=1}^{\theta} \left(\operatorname{Re} K_{\tau-1}^\theta(t) \cdot F_{\tau-1}^2(t, x) + \operatorname{Im} K_{\tau-1}^\theta(t) \cdot F_{\tau-1}^1(t, x) \right) - \operatorname{Im} C_\theta(t)$$

for all $(t, x) \in \tilde{\mathcal{T}} \times \mathcal{X}$, $\theta = 0, \dots, s_l - 1$,

where the functions

$$\operatorname{Re} K_{\tau-1}^\theta: t \rightarrow \int \sum_{j=1}^m \left(\binom{\theta}{\tau-1} \mu_{\theta-\tau+1, l}^{j\zeta} + \sum_{\delta=1}^{\theta-\tau} \binom{\theta}{\delta} (\mu_{\delta l}^{j\zeta} \operatorname{Re} K_{\tau-1}^{\theta-\delta}(t) - \tilde{\mu}_{\delta l}^{j\zeta} \operatorname{Im} K_{\tau-1}^{\theta-\delta}(t)) \right) dt_j,$$

$$\operatorname{Im} K_{\tau-1}^\theta: t \rightarrow \int \sum_{j=1}^m \left(\binom{\theta}{\tau-1} \tilde{\mu}_{\theta-\tau+1, l}^{j\zeta} + \sum_{\delta=1}^{\theta-\tau} \binom{\theta}{\delta} (\mu_{\delta l}^{j\zeta} \operatorname{Im} K_{\tau-1}^{\theta-\delta}(t) + \tilde{\mu}_{\delta l}^{j\zeta} \operatorname{Re} K_{\tau-1}^{\theta-\delta}(t)) \right) dt_j$$

$$\text{for all } t \in \tilde{\mathcal{T}}, \quad \tau = 1, \dots, \theta, \quad \theta = 1, \dots, s_l - 1,$$

$$\operatorname{Re} C_\theta: t \rightarrow \int \sum_{j=1}^m \left(\alpha_\theta(t, f_j(t)) + \sum_{\tau=1}^{\theta} \binom{\theta}{\tau} (\mu_{\tau l}^{j\zeta} \operatorname{Re} C_{\theta-\tau}(t) - \tilde{\mu}_{\tau l}^{j\zeta} \operatorname{Im} C_{\theta-\tau}(t)) \right) dt_j,$$

$$\operatorname{Im} C_\theta: t \rightarrow \int \sum_{j=1}^m \left(\beta_\theta(t, f_j(t)) + \sum_{\tau=1}^{\theta} \binom{\theta}{\tau} (\mu_{\tau l}^{j\zeta} \operatorname{Im} C_{\theta-\tau}(t) + \tilde{\mu}_{\tau l}^{j\zeta} \operatorname{Re} C_{\theta-\tau}(t)) \right) dt_j$$

$$\text{for all } t \in \tilde{\mathcal{T}}, \quad \theta = 0, \dots, s_l - 1, \quad \tilde{\mathcal{T}} \subset \mathcal{T},$$

the real numbers $\mu_{0l}^{*j\zeta} = \operatorname{Re} \lambda_l^j$, $\tilde{\mu}_{0l}^{j\zeta} = \operatorname{Im} \lambda_l^j$, $\mu_{\theta l}^{*j\zeta} = \operatorname{Re} \mu_{\theta l}^{j\zeta}$, $\tilde{\mu}_{\theta l}^{j\zeta} = \operatorname{Im} \mu_{\theta l}^{j\zeta}$, $j = 1, \dots, m$,
the vectors $\nu^{*\theta l} = \operatorname{Re} \nu^{\theta l}$, $\tilde{\nu}^{\theta l} = \operatorname{Im} \nu^{\theta l}$, $\theta = 0, \dots, s_l - 1$, and

$$\alpha_\theta(t, x) = \left(\nu^{*\theta l} x \cdot \cos \sum_{j=1}^m \tilde{\mu}_{0l}^{j\zeta} t_j + \tilde{\nu}^{\theta l} x \cdot \sin \sum_{j=1}^m \tilde{\mu}_{0l}^{j\zeta} t_j \right) \cdot \exp \left(- \sum_{j=1}^m \mu_{0l}^{*j\zeta} t_j \right),$$

$$\beta_\theta(t, x) = \left(\tilde{\nu}^{\theta l} x \cdot \cos \sum_{j=1}^m \tilde{\mu}_{0l}^{j\zeta}(t) t_j - \nu^{*\theta l} x \cdot \sin \sum_{j=1}^m \tilde{\mu}_{0l}^{j\zeta} t_j \right) \cdot \exp \left(- \sum_{j=1}^m \mu_{0l}^{*j\zeta} t_j \right)$$

$$\text{for all } (t, x) \in \tilde{\mathcal{T}} \times \mathbb{R}^n, \quad \theta = 0, \dots, s_l - 1.$$

Remark 2.4. Suppose the system (2.39) satisfies (2.2) and (2.40). Then 1-forms

$$\omega_{\tau-1}^\theta: t \rightarrow \sum_{j=1}^m \left(\binom{\theta}{\tau-1} \mu_{\theta-\tau+1, l}^{j\zeta} + \sum_{\delta=1}^{\theta-\tau} \binom{\theta}{\delta} \mu_{\delta l}^{j\zeta} \cdot K_{\tau-1}^{\theta-\delta}(t) \right) dt_j, \quad \tau = 1, \dots, \theta, \quad \theta = 1, \dots, s_l - 1,$$

$$\tilde{\omega}_\theta: t \rightarrow \sum_{j=1}^m \left(\nu^{\theta l} f_j(t) \cdot \varphi(t) + \sum_{\tau=1}^{\theta} \binom{\theta}{\tau} \mu_{\tau l}^{j\zeta} \cdot C_{\theta-\tau}(t) \right) dt_j \quad \text{for all } t \in \tilde{\mathcal{T}}, \quad \theta = 0, \dots, s_l - 1$$

are exact in a simply connected domain $\tilde{\mathcal{T}} \subset \mathcal{T}$.

3. First integrals of linear systems of ordinary differential equations

3.1. Linear homogeneous systems of ordinary differential equations

Let us consider a linear autonomous homogeneous system of ordinary differential equations

$$\frac{dx}{dt} = Ax, \quad (3.1)$$

where $x = \text{colon}(x_1, \dots, x_n) \in \mathbb{R}^n$, the $A = \|a_{ij}\|$ is a real $n \times n$ matrix.

Let B be the transpose of the matrix A .

Using methods of Section 2, we obtain the following statements.

Theorem 3.1. *Let $\nu \in \mathbb{C}^n$ be an eigenvector of the matrix B . Then the linear function $p: x \rightarrow \nu x$ for all $x \in \mathbb{R}^n$ is a partial integral of the system (3.1).*

3.1.1. Autonomous first integrals⁴

Theorem 3.2. *Let ν^1, ν^2 be real eigenvectors of the matrix B corresponding to the eigenvalues λ_1, λ_2 ($\lambda_1 \neq \lambda_2$), respectively. Then the system (3.1) has the first integral*

$$F: x \rightarrow |\nu^1 x|^{h_1} |\nu^2 x|^{h_2} \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset D(F),$$

where h_1, h_2 is a real solution to the equation $\lambda_1 h_1 + \lambda_2 h_2 = 0$ with $|h_1| + |h_2| \neq 0$.

Corollary 3.1. *If ν is a real eigenvector of the matrix B corresponding to the eigenvalue $\lambda = 0$, then the linear function $F: x \rightarrow \nu x$ for all $x \in \mathbb{R}^n$ is an autonomous first integral of the system of ordinary differential equations (3.1).*

Corollary 3.2. *Let $\lambda \neq 0$ be an eigenvalue of the matrix B corresponding to two real linearly independent eigenvectors ν^1, ν^2 . Then the system (3.1) has the autonomous first integral $F: x \rightarrow \frac{\nu^1 x}{\nu^2 x}$ for all $x \in \mathcal{X}$, where a domain $\mathcal{X} \subset \{x: \nu^2 x \neq 0\}$.*

For example, the autonomous system of ordinary differential equations

$$\begin{aligned} \frac{dx_1}{dt} &= x_1 - 2x_2 - x_4, & \frac{dx_2}{dt} &= -x_1 + 4x_2 - x_3 + 2x_4, \\ \frac{dx_3}{dt} &= 2x_2 + x_3 + x_4, & \frac{dx_4}{dt} &= 2x_1 - 4x_2 + 2x_3 - 2x_4 \end{aligned} \quad (3.2)$$

has the eigenvectors $\nu^1 = (1, -1, 1, -1), \nu^2 = (2, 2, 1, 1), \nu^3 = (1, 0, 1, 0), \nu^4 = (0, 2, 0, 1)$ corresponding to the eigenvalues $\lambda_1 = 0, \lambda_2 = \lambda_3 = 1, \lambda_4 = 2$, respectively. The functions

$$F_1: x \rightarrow x_1 - x_2 + x_3 - x_4 \quad \text{for all } x \in \mathbb{R}^4 \quad (\text{by Corollary 3.1}), \quad (3.3)$$

$$F_{23}: x \rightarrow \frac{2x_1 + 2x_2 + x_3 + x_4}{x_1 + x_3} \quad \text{for all } x \in \mathcal{X}_1 \quad (\text{by Corollary 3.2}), \quad (3.4)$$

$$F_{24}: x \rightarrow \frac{(2x_1 + 2x_2 + x_3 + x_4)^2}{2x_2 + x_4} \quad \text{for all } x \in \mathcal{X}_2 \quad (\text{by Theorem 3.2}), \quad (3.5)$$

where $\mathcal{X}_1 \subset \{x: x_1 + x_3 \neq 0\}, \mathcal{X}_2 \subset \{x: 2x_2 + x_4 \neq 0\}$, are autonomous first integrals of the system (3.2). The set of functionally independent first integrals F_1, F_{23}, F_{24} is a general autonomous integral of the system of ordinary differential equations (3.2).

⁴This Subsubsection has been published in *Vestnik of the Yanka Kupala Grodno State Univ.*, 2002, Ser. 2, No.2(11), 23 - 29; 2003, Ser. 2, No. 2(22), 50 - 60. Using method of Jordan canonical form, the same result was obtain by M. Falconi and J. Llibre in *Qualitative theory of dynamical systems*, 2004, No. 4, 233 - 254.

Theorem 3.3. Let $\nu = \nu^* + \tilde{\nu} i$ ($\nu^* = \operatorname{Re} \nu$, $\tilde{\nu} = \operatorname{Im} \nu$) be an eigenvector of the matrix B corresponding to the complex eigenvalue $\lambda = \lambda^* + \tilde{\lambda} i$ ($\lambda^* = \operatorname{Re} \lambda$, $\tilde{\lambda} = \operatorname{Im} \lambda \neq 0$). Then the system of ordinary differential equations (3.1) has the autonomous first integral

$$F: x \rightarrow ((\nu^* x)^2 + (\tilde{\nu} x)^2) \cdot \exp\left(-2 \frac{\lambda^*}{\tilde{\lambda}} \operatorname{arctg} \frac{\tilde{\nu} x}{\nu^* x}\right) \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^* x \neq 0\}$.

Theorem 3.4. Let $\nu^1 = \nu^{*1} + \tilde{\nu}^1 i$ ($\nu^{*1} = \operatorname{Re} \nu^1$, $\tilde{\nu}^1 = \operatorname{Im} \nu^1$) be an eigenvector of the matrix B corresponding to the complex eigenvalue $\lambda_1 = \lambda_1^* + \tilde{\lambda}_1 i$ ($\lambda_1^* = \operatorname{Re} \lambda_1$, $\tilde{\lambda}_1 = \operatorname{Im} \lambda_1 \neq 0$), ν^2 be a real eigenvector of the matrix B corresponding to the eigenvalue $\lambda_2 \neq 0$. Then the system of ordinary differential equations (3.1) has the autonomous first integral

$$F: x \rightarrow \nu^2 x \cdot \exp\left(-\frac{\lambda_2}{\tilde{\lambda}_1} \operatorname{arctg} \frac{\tilde{\nu}^1 x}{\nu^{*1} x}\right) \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^{*1} x \neq 0\}$.

The autonomous linear system of ordinary differential equations

$$\frac{dx_1}{dt} = 2x_1 + x_2, \quad \frac{dx_2}{dt} = x_1 + 3x_2 - x_3, \quad \frac{dx_3}{dt} = -x_1 + 2x_2 + 3x_3 \quad (3.6)$$

has the eigenvalues $\lambda_1 = 3 + i$, $\lambda_2 = 2$ corresponding to two eigenvectors $\nu^1 = (1, i, -1)$, $\nu^2 = (3, -1, -1)$, respectively, and the first autonomous integrals

$$F_1: x \rightarrow ((x_1 - x_3)^2 + x_2^2) \exp\left(-6 \operatorname{arctg} \frac{x_2}{x_1 - x_3}\right) \quad \text{for all } x \in \mathcal{X} \quad (\text{by Theorem 3.3}), \quad (3.7)$$

$$F_2: x \rightarrow (3x_1 - x_2 - x_3) \exp\left(-2 \operatorname{arctg} \frac{x_2}{x_1 - x_3}\right) \quad \text{for all } x \in \mathcal{X} \quad (\text{by Theorem 3.4}), \quad (3.8)$$

where a domain $\mathcal{X} \subset \{x: x_1 - x_3 \neq 0\}$.

Theorem 3.5. Let $\nu^1 = \nu^{*1} + \tilde{\nu}^1 i$ and $\nu^2 = \nu^{*2} + \tilde{\nu}^2 i$ be two eigenvectors of the matrix B corresponding to the complex eigenvalues $\lambda_1 = \lambda_1^* + \tilde{\lambda}_1 i$ and $\lambda_2 = \lambda_2^* + \tilde{\lambda}_2 i$ ($\lambda_1 \neq \bar{\lambda}_2$), respectively. Then the system (3.1) has the autonomous first integral

$$F: x \rightarrow \tilde{\lambda}_1 \operatorname{arctg} \frac{\tilde{\nu}^2 x}{\nu^{*2} x} - \tilde{\lambda}_2 \operatorname{arctg} \frac{\tilde{\nu}^1 x}{\nu^{*1} x} \quad \text{for all } x \in \mathcal{X},$$

where the vectors $\nu^\tau = \nu^{*\tau} + \tilde{\nu}^\tau i$, $\nu^{*\tau} = \operatorname{Re} \nu^\tau$, $\tilde{\nu}^\tau = \operatorname{Im} \nu^\tau$, the numbers $\lambda_\tau = \lambda_\tau^* + \tilde{\lambda}_\tau i$, $\lambda_\tau^* = \operatorname{Re} \lambda_\tau$, $\tilde{\lambda}_\tau = \operatorname{Im} \lambda_\tau \neq 0$, $\tau = 1, 2$, a domain $\mathcal{X} \subset \{x: \nu^{*2} x \neq 0 \wedge \nu^{*1} x \neq 0\}$.

As an example, the linear autonomous system of ordinary differential equations

$$\begin{aligned} \frac{dx_1}{dt} &= -3x_1 + x_2 + 4x_3 + 2x_4, & \frac{dx_2}{dt} &= 8x_1 - 3x_2 - 2x_3 + 6x_4, \\ \frac{dx_3}{dt} &= -9x_1 + 3x_2 + 4x_3 - 4x_4, & \frac{dx_4}{dt} &= 6x_1 - 3x_2 - 4x_3 + 2x_4 \end{aligned} \quad (3.9)$$

has the eigenvalues $\lambda_1 = i$, $\lambda_2 = 2i$ corresponding to the eigenvectors $\nu^1 = (1 - i, -1 + 2i, 2i, 2)$, $\nu^2 = (i, -1, i, 1 + 2i)$, respectively. The functionally independent first integrals

$$F_1: x \rightarrow (x_1 - x_2 + 2x_4)^2 + (-x_1 + 2x_2 + 2x_3)^2 \quad \text{for all } x \in \mathbb{R}^4 \quad (\text{by Theorem 3.3}), \quad (3.10)$$

$$F_2: x \rightarrow (-x_2 + x_4)^2 + (x_1 + x_3 + 2x_4)^2 \quad \text{for all } x \in \mathbb{R}^4 \quad (\text{by Theorem 3.3}), \quad (3.11)$$

and (by Theorem 3.5)

$$F_3: x \rightarrow \operatorname{arctg} \frac{x_1 + x_3 + 2x_4}{-x_2 + x_4} - 2 \operatorname{arctg} \frac{-x_1 + 2x_2 + 2x_3}{x_1 - x_2 + 2x_4} \quad \text{for all } x \in \mathcal{X} \quad (3.12)$$

are a general autonomous integral on a domain $\mathcal{X} \subset \{x: x_1 - x_2 + 2x_4 \neq 0 \wedge x_2 - x_4 \neq 0\}$ of the linear system of ordinary differential equations (3.9).

Definition 3.1. Let ν^0 be an eigenvector of the matrix B corresponding to the eigenvalue λ with elementary divisor of multiplicity m . A non-zero vector $\nu^k \in \mathbb{C}^n$ is called a **generalized eigenvector of order k** for λ if and only if

$$(B - \lambda E) \nu^k = k \cdot \nu^{k-1}, \quad k = 1, \dots, m-1,$$

where E is the $n \times n$ identity matrix.

Theorem 3.6. Let λ be an eigenvalue of the matrix B with the elementary divisor of multiplicity m ($m \geq 2$) corresponding to the real eigenvector ν^0 and to the real order 1 generalized eigenvector ν^1 . Then the system (3.1) has the autonomous first integral

$$F: x \rightarrow \nu^0 x \exp\left(-\lambda \frac{\nu^1 x}{\nu^0 x}\right) \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^0 x \neq 0\}$.

Corollary 3.3. Let $\lambda = \lambda^* + \tilde{\lambda}i$ ($\lambda^* = \operatorname{Re} \lambda$, $\tilde{\lambda} = \operatorname{Im} \lambda \neq 0$) be a complex eigenvalue of the matrix B with the elementary divisor of multiplicity m ($m \geq 2$) corresponding to the eigenvector $\nu^0 = \nu^{*0} + \tilde{\nu}^0 i$ and to the order 1 generalized eigenvector $\nu^1 = \nu^{*1} + \tilde{\nu}^1 i$. Then the system (3.1) has the autonomous first integrals

$$F_1: x \rightarrow \left((\nu^{*0} x)^2 + (\tilde{\nu}^0 x)^2\right) \exp\left(-2 \frac{\lambda^* \alpha(x) - \tilde{\lambda} \beta(x)}{(\nu^{*0} x)^2 + (\tilde{\nu}^0 x)^2}\right) \quad \text{for all } x \in \mathcal{X}$$

and

$$F_2: x \rightarrow \operatorname{arctg} \frac{\tilde{\nu}^0 x}{\nu^{*0} x} - \frac{\tilde{\lambda} \alpha(x) + \lambda^* \beta(x)}{(\nu^{*0} x)^2 + (\tilde{\nu}^0 x)^2} \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^{*0} x \neq 0\}$, the vectors $\nu^\tau = \operatorname{Re} \nu^\tau$, $\tilde{\nu}^\tau = \operatorname{Im} \nu^\tau$, $\tau = 0, 1$, the polynomials $\alpha: x \rightarrow \nu^{*0} x \nu^{*1} x + \tilde{\nu}^0 x \tilde{\nu}^1 x$, $\beta: x \rightarrow \nu^{*0} x \tilde{\nu}^1 x - \tilde{\nu}^0 x \nu^{*1} x$ for all $x \in \mathbb{R}^n$.

Corollary 3.4. Let $\lambda_1 = 0$ be an eigenvalue with the elementary divisor of multiplicity m ($m \geq 2$) of the matrix B corresponding to the real eigenvector ν^0 and the real order 1 generalized eigenvector ν^1 . Let λ_2 be an eigenvalue of the matrix B corresponding to the real eigenvector ν^2 . Then the system (3.1) has the autonomous first integrals

$$F: x \rightarrow \nu^2 x \exp\left(-\lambda_2 \frac{\nu^1 x}{\nu^0 x}\right) \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^0 x \neq 0\}$.

The linear autonomous system of ordinary differential equations

$$\frac{dx_1}{dt} = 4x_1 - 5x_2 + 2x_3, \quad \frac{dx_2}{dt} = 5x_1 - 7x_2 + 3x_3, \quad \frac{dx_3}{dt} = 6x_1 - 9x_2 + 4x_3 \quad (3.13)$$

has the eigenvalue $\lambda_1 = 0$ with the elementary divisor λ^2 of multiplicity 2 corresponding to the eigenvector $\nu^0 = (1, -2, 1)$ and the order 1 generalized eigenvector $\nu^1 = (0, -1, 1)$, and the simple eigenvalue $\lambda_3 = 1$ with the elementary divisor $\lambda - 1$ corresponding to the eigenvector $\nu^3 = (3, -3, 1)$. The functionally independent functions

$$F_1: x \rightarrow x_1 - 2x_2 + x_3 \quad \text{for all } x \in \mathbb{R}^3 \quad (\text{by Theorem 3.6}) \quad (3.14)$$

and (by Corollary 3.4)

$$F_2: x \rightarrow (3x_1 - 3x_2 + x_3) \exp \frac{x_2 - x_3}{x_1 - 2x_2 + x_3} \quad \text{for all } x \in \mathcal{X}, \quad (3.15)$$

where $\mathcal{X} \subset \{x: x_1 - 2x_2 + x_3 \neq 0\}$, are autonomous first integrals of the system (3.13).

Corollary 3.5. *Let $\lambda_1 = 0$ be an eigenvalue with elementary divisor of multiplicity m ($m \geq 2$) of the matrix B corresponding to the real eigenvector ν^0 and to the real order 1 generalized eigenvector ν^1 . Let $\lambda_2 = \lambda_2^* + \tilde{\lambda}_2 i$ ($\lambda_2^* = \operatorname{Re} \lambda_2$, $\tilde{\lambda}_2 = \operatorname{Im} \lambda_2 \neq 0$) be a complex eigenvalue of the matrix B corresponding to the eigenvector $\nu^2 = \tilde{\nu}^2 + \tilde{\nu}^2 i$. Then the system of differential equations (3.1) has two autonomous first integrals*

$$F_1: x \rightarrow \left((\tilde{\nu}^2 x)^2 + (\tilde{\nu}^2 x)^2 \right) \exp \left(-2 \lambda_2^* \frac{\nu^1 x}{\nu^0 x} \right) \quad \text{for all } x \in \mathcal{X}$$

and

$$F_2: x \rightarrow \operatorname{arctg} \frac{\tilde{\nu}^2 x}{\nu^2 x} - \tilde{\lambda}_2 \frac{\nu^1 x}{\nu^0 x} \quad \text{for all } x \in \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \nu^0 x \neq 0, \tilde{\nu}^2 x \neq 0\}$.

Theorem 3.7. *Let λ be an eigenvalue with elementary divisor of multiplicity $m \geq 2$ of the matrix B corresponding to the real eigenvector ν^0 and to the real generalized eigenvectors ν^k , $k = \overline{1, m-1}$. Then the system (3.1) has the first integrals*

$$F_g: x \rightarrow \Psi_g(x) \quad \text{for all } x \in \mathcal{X}, \quad g = 2, \dots, m-1, \quad (3.16)$$

where the functions $\Psi_g: \mathcal{X} \rightarrow \mathbb{R}$ is the solution to system

$$\nu^k x = \sum_{i=1}^k \binom{k-1}{i-1} \Psi_i(x) \nu^{k-i} x, \quad k = 1, \dots, m-1, \quad \mathcal{X} \subset \{x: \nu^0 x \neq 0\}.$$

Remark 3.1. Let $\lambda = \lambda^* + \tilde{\lambda} i$ ($\lambda^* = \operatorname{Re} \lambda$, $\tilde{\lambda} = \operatorname{Im} \lambda \neq 0$) be a complex eigenvalue with elementary divisor of multiplicity m ($m \geq 2$) of the matrix B corresponding to the complex eigenvector $\nu^0 = \tilde{\nu}^0 + \tilde{\nu}^0 i$. Taking into account (3.16), we obtain the real-valued first integrals of the system of ordinary differential equations (3.1):

$$F_{g,1}: x \rightarrow \operatorname{Re} \Psi_g(x), \quad F_{g,2}: x \rightarrow \operatorname{Im} \Psi_g(x) \quad \text{for all } x \in \mathcal{X}, \quad g = 2, \dots, m-1,$$

where a domain $\mathcal{X} \subset \{x: (\tilde{\nu}^0 x)^2 + (\tilde{\nu}^0 x)^2 \neq 0\}$.

For example, the system of ordinary differential equations

$$\frac{dx_1}{dt} = 4x_1 - x_2, \quad \frac{dx_2}{dt} = 3x_1 + x_2 - x_3, \quad \frac{dx_3}{dt} = x_1 + x_3 \quad (3.17)$$

has the eigenvalue $\lambda_1 = 2$ with the elementary divisor $(\lambda - 2)^3$ of multiplicity 3 corresponding to the eigenvector $\nu^0 = (1, -1, 1)$ and the generalized eigenvectors $\nu^1 = (1, 0, -1)$, $\nu^2 = (0, 0, 2)$. First integrals of the system of differential equations (3.17) are the functions

$$F_1: (x_1, x_2, x_3) \rightarrow (x_1 - x_2 + x_3) \exp \left(-2 \frac{x_1 - x_3}{x_1 - x_2 + x_3} \right) \quad (\text{by Theorem 3.6}) \quad (3.18)$$

and (by Theorem 3.7)

$$F_2: (x_1, x_2, x_3) \rightarrow \frac{(x_1 - x_3)^2 - 2x_3(x_1 - x_2 + x_3)}{(x_1 - x_2 + x_3)^2} \quad \text{for all } (x_1, x_2, x_3) \in \mathcal{X}, \quad (3.19)$$

where a domain $\mathcal{X} \subset \{(x_1, x_2, x_3): x_1 - x_2 + x_3 \neq 0\}$.

The autonomous system of ordinary differential equations

$$\begin{aligned} \frac{dx_1}{dt} &= x_1 - 2x_2 + x_3 - 2x_6, & \frac{dx_2}{dt} &= 3x_2 - x_3 - x_5 + 2x_6, \\ \frac{dx_3}{dt} &= -x_1 + x_3 + 2x_4 + 2x_5, & \frac{dx_4}{dt} &= -x_1 + x_4 + x_5 + x_6, \\ \frac{dx_5}{dt} &= x_1 + x_2 + x_5, & \frac{dx_6}{dt} &= x_1 - x_2 + x_3 - x_4 - x_6 \end{aligned} \quad (3.20)$$

has the complex eigenvalue $\lambda_1 = 1+i$ with the elementary divisor $(\lambda-1-i)^3$ corresponding to the eigenvector $\nu^0 = (1, 1, 0, 0, i, 0)$ and to the generalized eigenvectors $\nu^1 = (0, 1, 0, i, i, 1)$, $\nu^2 = (0, 1, i, 0, i, 0)$. A general autonomous integral of the system (3.20) is the functions

$$F_1: x \rightarrow P(x) \exp(-2\varphi(x)) \quad \text{for all } x \in \mathcal{X} \quad (\text{by Theorem 3.3}), \quad (3.21)$$

$$F_2: x \rightarrow P(x) \exp\left(-2 \frac{\alpha(x) - \beta(x)}{P(x)}\right) \quad \text{for all } x \in \mathcal{X} \quad (\text{by Corollary 3.3}), \quad (3.22)$$

$$F_3: x \rightarrow \varphi(x) - \frac{\alpha(x) + \beta(x)}{P(x)} \quad \text{for all } x \in \mathcal{X} \quad (\text{by Corollary 3.3}), \quad (3.23)$$

$$F_4: x \rightarrow \frac{\gamma(x)P(x) + \beta^2(x) - \alpha^2(x)}{P^2(x)} \quad \text{for all } x \in \mathcal{X} \quad (\text{by Theorem 3.7}), \quad (3.24)$$

and

$$F_5: x \rightarrow \frac{\delta(x)P(x) - 2\alpha(x)\beta(x)}{P^2(x)} \quad \text{for all } x \in \mathcal{X} \quad (\text{by Theorem 3.7}), \quad (3.25)$$

where

$$P: x \rightarrow (x_1 + x_2)^2 + x_5^2, \quad \alpha: x \rightarrow (x_1 + x_2)(x_2 + x_6) + x_5(x_4 + x_5),$$

$$\beta: x \rightarrow (x_1 + x_2)(x_4 + x_5) - x_5(x_2 + x_6), \quad \gamma: x \rightarrow x_2(x_1 + x_2) + x_5(x_3 + x_5),$$

$$\delta: x \rightarrow (x_1 + x_2)(x_3 + x_5) - x_2x_5 \quad \text{for all } x \in \mathbb{R}^6,$$

$$\varphi: x \rightarrow \arctg \frac{x_5}{x_1 + x_2} \quad \text{for all } x \in \mathcal{X}, \quad \mathcal{X} \subset \{x: x_1 + x_2 \neq 0\}.$$

3.1.2. Nonautonomous first integrals

Theorem 3.8. *Let ν be a real eigenvector of the matrix B corresponding to the eigenvalue λ . Then the system (3.1) has the first integral*

$$F: (t, x) \rightarrow \nu x \exp(-\lambda t) \quad \text{for all } (t, x) \in \mathbb{R}^{n+1}.$$

For example, the four dimensional system (3.2) has the eigenvalue $\lambda_2 = 1$ corresponding to the eigenvector $\nu^2 = (2, 2, 1, 1)$ and the first integral

$$F: (t, x) \rightarrow (2x_1 + 2x_2 + x_3 + x_4) e^{-t} \quad \text{for all } (t, x) \in \mathbb{R}^5 \quad (\text{by Theorem 3.8}).$$

The first integrals (3.3), (3.4), (3.5), and F are a general integral on a domain $\mathbb{R} \times \mathcal{X}$ of the system (3.2), where $\mathcal{X} \subset \{x: x_1 + x_3 \neq 0 \wedge 2x_2 + x_4 \neq 0\} \subset \mathbb{R}^4$.

Using the eigenvector $\nu^2 = (3, -1, -1)$ corresponding to the eigenvalue $\lambda_2 = 2$, we can build the first integral of the linear system (3.6):

$$F: (t, x_1, x_2, x_3) \rightarrow (3x_1 - x_2 - x_3)e^{-2t} \quad \text{for all } (t, x_1, x_2, x_3) \in \mathbb{R}^4 \quad (\text{by Theorem 3.8}).$$

The functionally independent first integrals (3.7), (3.8), and F are a general integral on a domain $\mathbb{R} \times \mathcal{X}$ of the system (3.6), where a domain $\mathcal{X} \subset \{(x_1, x_2, x_3): x_1 - x_3 \neq 0\}$.

Corollary 3.6. *Let $\nu = \overset{*}{\nu} + \tilde{\nu}i$ ($\overset{*}{\nu} = \text{Re } \nu$, $\tilde{\nu} = \text{Im } \nu$) be an eigenvector of the matrix B corresponding to the complex eigenvalue $\lambda = \overset{*}{\lambda} + \tilde{\lambda}i$ ($\overset{*}{\lambda} = \text{Re } \lambda$, $\tilde{\lambda} = \text{Im } \lambda \neq 0$). Then the system of ordinary differential equations (3.1) has the first integrals*

$$F_1: (t, x) \rightarrow \left((\overset{*}{\nu}x)^2 + (\tilde{\nu}x)^2 \right) \exp(-2\overset{*}{\lambda}t) \quad \text{for all } (t, x) \in \mathbb{R}^{n+1}$$

and

$$F_2: (t, x) \rightarrow \arctg \frac{\tilde{\nu}x}{\overset{*}{\nu}x} - \tilde{\lambda}t \quad \text{for all } (t, x) \in \mathbb{R} \times \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: \overset{*}{\nu}x \neq 0\}$.

As an example, the system of linear differential equations (3.9) has the eigenvalue $\lambda_1 = i$ corresponding to the eigenvector $\nu^1 = (1 - i, -1 + 2i, 2i, 2)$ and the first integral

$$F: (t, x) \rightarrow \arctg \frac{-x_1 + 2x_2 + 2x_3}{x_1 - x_2 + 2x_4} - t \quad \text{for all } (t, x) \in \mathbb{R} \times \mathcal{X}_1 \quad (\text{by Corollary 3.6}),$$

where a domain $\mathcal{X}_1 \subset \{x: x_1 - x_2 + 2x_4 \neq 0\}$.

The first integrals (3.10), (3.11), (3.12) and F are a general integral on a domain $\mathbb{R} \times \mathcal{X}$ of the system (3.9), where a domain $\mathcal{X} \subset \{x: x_1 - x_2 + 2x_4 \neq 0 \wedge x_4 - x_2 \neq 0\} \subset \mathbb{R}^4$.

Theorem 3.9. *Let λ be an eigenvalue of the matrix B with the elementary divisor of multiplicity m ($m \geq 2$) corresponding to the real eigenvector ν^0 and the real order 1 generalized eigenvector ν^1 . Then the system (3.1) has the first integral*

$$F: (t, x) \rightarrow \frac{\nu^1 x}{\nu^0 x} - t \quad \text{for all } (t, x) \in \mathbb{R} \times \mathcal{X}, \quad (3.26)$$

where a domain $\mathcal{X} \subset \{x: \nu^0 x \neq 0\}$.

The system of ordinary differential equations (3.13) has the eigenvalue $\lambda_1 = 0$ corresponding to the eigenvector $\nu^0 = (1, -2, 1)$ and to the order 1 generalized eigenvector $\nu^1 = (0, -1, 1)$. From Theorem 3.9 it follows that the function

$$F: (t, x_1, x_2, x_3) \rightarrow \frac{x_3 - x_2}{x_1 - 2x_2 + x_3} - t \quad \text{for all } (t, x_1, x_2, x_3) \in \mathbb{R} \times \mathcal{X}$$

is a first integral of the system (3.13), where a domain $\mathcal{X} \subset \{(x_1, x_2, x_3): x_1 - 2x_2 + x_3 \neq 0\}$.

A general integral on a domain $\mathbb{R} \times \mathcal{X}$ of the system of ordinary differential equations (3.13) is the functionally independent first integrals (3.14), (3.15), and F .

Consider the system (3.17). Using the eigenvalue $\lambda_1 = 2$ corresponding to the eigenvector $\nu^0 = (1, -1, 1)$ and to the order 1 generalized eigenvector $\nu^1 = (1, 0, -1)$, we can build the first integral of the system of ordinary differential equations (3.17):

$$F: (t, x_1, x_2, x_3) \rightarrow \frac{x_1 - x_3}{x_1 - x_2 + x_3} - t \quad \text{for all } (t, x_1, x_2, x_3) \in \mathbb{R} \times \mathcal{X} \quad (\text{by Theorem 3.9}).$$

The first integrals (3.18), (3.19), and F are a general integral on a domain $\mathbb{R} \times \mathcal{X}$ of the system (3.17), where a domain $\mathcal{X} \subset \{(x_1, x_2, x_3): x_1 - x_2 + x_3 \neq 0\} \subset \mathbb{R}^3$.

Remark 3.2. Suppose $\lambda = \overset{*}{\lambda} + \tilde{\lambda} i$ ($\overset{*}{\lambda} = \operatorname{Re} \lambda$, $\tilde{\lambda} = \operatorname{Im} \lambda \neq 0$) is a complex eigenvalue of the matrix B corresponding to the eigenvector $\nu^0 = \overset{*}{\nu}^0 + \tilde{\nu}^0 i$ and to the generalised eigenvector $\nu^1 = \overset{*}{\nu}^1 + \tilde{\nu}^1 i$. Using (3.26), we get the real-valued first integrals of the system (3.1):

$$F_1: (t, x) \rightarrow \frac{\overset{*}{\nu}^0 x \overset{*}{\nu}^1 x + \tilde{\nu}^0 x \tilde{\nu}^1 x}{(\overset{*}{\nu}^0 x)^2 + (\tilde{\nu}^0 x)^2} - t, \quad F_2: (t, x) \rightarrow \frac{\overset{*}{\nu}^0 x \tilde{\nu}^1 x - \tilde{\nu}^0 x \overset{*}{\nu}^1 x}{(\overset{*}{\nu}^0 x)^2 + (\tilde{\nu}^0 x)^2} \quad \text{for all } (t, x) \in \mathbb{R} \times \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: (\overset{*}{\nu}^0 x)^2 + (\tilde{\nu}^0 x)^2 \neq 0\}$.

Consider the system (3.20). Using the eigenvalue $\lambda_1 = 1 + i$ corresponding to the eigenvector $\nu^0 = (1, 1, 0, 0, i, 0)$ and to the order 1 generalized eigenvector $\nu^1 = (0, 1, 0, i, i, 1)$, we can build the first integral of the system of ordinary differential equations (3.20):

$$F: (t, x) \rightarrow \frac{(x_1 + x_2)(x_2 + x_6) + x_5(x_4 + x_5)}{(x_1 + x_2)^2 + x_5^2} - t \quad \text{for all } (t, x) \in \mathbb{R} \times \mathcal{X},$$

where a domain $\mathcal{X} \subset \{x: x_1 + x_2 \neq 0\} \subset \mathbb{R}^6$.

The functionally independent first integrals (3.21), (3.22), (3.23), (3.24), (3.25), and F are a general integral of the system of ordinary differential equations (3.20).

3.2. Linear nonhomogeneous systems of ordinary differential equations

We consider a system of n first order constant-coefficient linear nonhomogeneous ordinary differential equations

$$\frac{dx}{dt} = Ax + f(t), \quad (3.27)$$

where $x = \operatorname{colon}(x_1, \dots, x_n) \in \mathbb{R}^n$, $A = \|a_{ij}\|$ is a real $n \times n$ matrix, and the vector function $f: t \rightarrow \operatorname{colon}(f_1(t), \dots, f_n(t))$ for all $t \in J$ is continuous on an interval $J \subset \mathbb{R}$.

Let B be the transpose of the matrix A .

Theorem 3.10. Suppose ν is a real eigenvector of the matrix B corresponding to the eigenvalue λ . Then a first integral of the system (3.27) is the function

$$F: (t, x) \rightarrow \nu x \cdot \exp(-\lambda t) - \int \nu f(t) \cdot \exp(-\lambda t) dt \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad \tilde{J} \subset J.$$

Corollary 3.7. Let $\nu = \overset{*}{\nu} + \tilde{\nu} i$ ($\overset{*}{\nu} = \operatorname{Re} \nu$, $\tilde{\nu} = \operatorname{Im} \nu$) be an eigenvector of the matrix B corresponding to the complex eigenvalue $\lambda = \overset{*}{\lambda} + \tilde{\lambda} i$ ($\overset{*}{\lambda} = \operatorname{Re} \lambda$, $\tilde{\lambda} = \operatorname{Im} \lambda \neq 0$). Then first integrals of the system of ordinary differential equations (3.27) are the functions

$$F_\tau: (t, x) \rightarrow \alpha_\tau(t, x) - \int \alpha_\tau(t, f(t)) dt \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad \tau = 1, 2, \quad \tilde{J} \subset J,$$

where the functions $\alpha_1(t, x) = (\overset{*}{\nu} x \cdot \cos \tilde{\lambda} t + \tilde{\nu} x \cdot \sin \tilde{\lambda} t) \cdot \exp(-\overset{*}{\lambda} t)$ for all $(t, x) \in J \times \mathbb{R}^n$, $\alpha_2(t, x) = (\tilde{\nu} x \cdot \cos \tilde{\lambda} t - \overset{*}{\nu} x \cdot \sin \tilde{\lambda} t) \cdot \exp(-\overset{*}{\lambda} t)$ for all $(t, x) \in J \times \mathbb{R}^n$.

Remark 3.3. Under the conditions of Corollary 3.7, we have the function

$$F: (t, x) \rightarrow \left((\overset{*}{\nu} x)^2 + (\tilde{\nu} x)^2 \right) \exp(-2 \overset{*}{\lambda} t) - 2 \left(\alpha_1(t, x) \int \alpha_1(t, f(t)) dt + \alpha_2(t, x) \int \alpha_2(t, f(t)) dt \right) + \\ + \left(\int \alpha_1(t, f(t)) dt \right)^2 + \left(\int \alpha_2(t, f(t)) dt \right)^2 \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n$$

is also a first integral of the system (3.27).

For example, the linear nonhomogeneous system of ordinary differential equations

$$\frac{dx_1}{dt} = 2x_1 + x_2 + 2e^{2t}, \quad \frac{dx_2}{dt} = x_1 + 3x_2 - x_3 + 10, \quad \frac{dx_3}{dt} = -x_1 + 2x_2 + 3x_3 + e^{3t}$$

has the eigenvalue $\lambda_1 = 2$ corresponding to the eigenvector $\nu^1 = (3, -1, -1)$, the eigenvalue $\lambda_2 = 3 + i$ corresponding to the eigenvector $\nu^2 = (1, i, -1)$, and the first integrals

$$\begin{aligned} F_1 &: (t, x_1, x_2, x_3) \rightarrow (3x_1 - x_2 - x_3 - 5)e^{-2t} + e^t - 6t \quad (\text{by Theorem 3.10}), \\ F_2 &: (t, x_1, x_2, x_3) \rightarrow ((x_1 - x_3 + 1) \cos t + (x_2 + 3) \sin t)e^{-3t} + (\cos t - \sin t)e^{-t} + \sin t, \\ F_3 &: (t, x_1, x_2, x_3) \rightarrow ((x_2 + 3) \cos t + (x_3 - x_1 - 1) \sin t)e^{-3t} - (\cos t + \sin t)e^{-t} + \cos t \\ &\quad \text{for all } (t, x_1, x_2, x_3) \in \mathbb{R}^4 \quad (\text{by Corollary 3.7}). \end{aligned}$$

Theorem 3.11. *Let λ be an eigenvalue with the elementary divisor of multiplicity m ($m \geq 2$) of the matrix B corresponding to the real eigenvector ν^0 and to the real generalized eigenvectors ν^k , $k = 1, \dots, m-1$. Then the system (3.27) has the first integrals*

$$\begin{aligned} F_{k+1} &: (t, x) \rightarrow (\nu^k x) \cdot \exp(-\lambda t) - \sum_{i=0}^{k-1} \binom{k}{i} t^{k-i} \cdot F_{i+1}(t, x) - \\ &\quad - \int \left((\nu^k f(t)) \cdot \exp(-\lambda t) + k \cdot \int \left((\nu^{k-1} f(t)) \cdot \exp(-\lambda t) + \right. \right. \\ &\quad \left. \left. + (k-1) \cdot \int \left((\nu^{k-2} f(t)) \cdot \exp(-\lambda t) + \dots + 2 \int \left((\nu^1 f(t)) \cdot \exp(-\lambda t) + \right. \right. \right. \\ &\quad \left. \left. \left. + \int (\nu^0 f(t)) \cdot \exp(-\lambda t) dt \right) dt \dots \right) dt \right) dt \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad k = 1, \dots, m-1, \end{aligned}$$

where the first integral (by Theorem 3.10)

$$F_1: (t, x) \rightarrow (\nu^0 x) \cdot \exp(-\lambda t) - \int (\nu^0 f(t)) \cdot \exp(-\lambda t) dt \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad \tilde{J} \subset J.$$

As an example, the system of ordinary differential equations

$$\frac{dx_1}{dt} = 4x_1 - x_2 + e^{3t}, \quad \frac{dx_2}{dt} = 3x_1 + x_2 - x_3 + 8t, \quad \frac{dx_3}{dt} = x_1 + x_3 + 4$$

has the eigenvalue $\lambda_1 = 2$ with the elementary divisor $(\lambda - 2)^3$ of multiplicity 3 corresponding to the eigenvector $\nu^0 = (1, -1, 1)$ and the generalized eigenvectors $\nu^1 = (1, 0, -1)$, $\nu^2 = (-2, 2, 0)$. First integrals of this system of differential equations are the functions

$$\begin{aligned} F_1 &: (t, x) \rightarrow (x_1 - x_2 + x_3 - 4t)e^{-2t} - e^t \quad \text{for all } (t, x) \in \mathbb{R}^4 \quad (\text{by Theorem 3.10}), \\ F_2 &: (t, x) \rightarrow (x_1 - x_3 + 2t - 1)e^{-2t} - t F_1(t, x) - 2e^t \quad \text{for all } (t, x) \in \mathbb{R}^4 \quad (\text{by Theorem 3.11}), \\ &\quad \text{and (by Theorem 3.11)} \end{aligned}$$

$$F_3: (t, x) \rightarrow 2(x_2 - x_1 + 3t + 2)e^{-2t} - t^2 F_1(t, x) - 2t F_2(t, x) - 2e^t \quad \text{for all } (t, x) \in \mathbb{R}^4.$$

Theorem 3.12. *Let $\lambda = \lambda^* + \tilde{\lambda} i$ ($\tilde{\lambda} \neq 0$) be an eigenvalue with the elementary divisor of multiplicity m ($m \geq 2$) of the matrix B corresponding to the eigenvector $\nu^0 = \tilde{\nu}^0 + \tilde{\nu}^0 i$ and to the generalized eigenvectors $\nu^k = \tilde{\nu}^k + \tilde{\nu}^k i$, $k = 1, \dots, m-1$. Then the system of ordinary differential equations (3.27) has the first integrals*

$$\begin{aligned}
F_{\tau,k+1} : (t, x) &\rightarrow \alpha_{\tau k}(t, x) - \sum_{\xi=0}^{k-1} \binom{k}{\xi} t^{k-\xi} \cdot F_{\tau,\xi+1}(t, x) - \\
&- \int \left(\alpha_{\tau k}(t, f(t)) + k \cdot \int \left(\alpha_{\tau,k-1}(t, f(t)) + (k-1) \cdot \int \left(\alpha_{\tau,k-2}(t, f(t)) + \right. \right. \right. \\
&\quad \left. \left. \left. + \dots + 2 \int \left(\alpha_{\tau 1}(t, f(t)) + \int \alpha_{\tau 0}(t, f(t)) dt \right) dt \dots \right) dt \right) dt \\
&\text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad k = 1, \dots, m-1, \quad \tau = 1, 2, \quad \tilde{J} \subset J,
\end{aligned}$$

where the functions

$$\alpha_{1k}(t, x) = (\tilde{\nu}^k x \cdot \cos \tilde{\lambda} t + \tilde{\nu}^k x \cdot \sin \tilde{\lambda} t) \cdot \exp(-\tilde{\lambda}^* t) \quad \text{for all } (t, x) \in \mathbb{R}^{n+1}, \quad k = 0, \dots, m-1,$$

$$\alpha_{2k}(t, x) = (\tilde{\nu} x^k \cdot \cos \tilde{\lambda} t - \tilde{\nu}^* x \cdot \sin \tilde{\lambda} t) \cdot \exp(-\tilde{\lambda}^* t) \quad \text{for all } (t, x) \in \mathbb{R}^{n+1}, \quad k = 0, \dots, m-1,$$

and the first integrals (by Corollary 3.7)

$$F_{\tau 1} : (t, x) \rightarrow \alpha_{\tau 0}(t, x) - \int \alpha_{\tau 0}(t, f(t)) dt \quad \text{for all } (t, x) \in \tilde{J} \times \mathbb{R}^n, \quad \tau = 1, 2.$$

References

- [1] B.V. Shabat, *An introduction to complex analysis: functions of several variables* (Russian), P. 2, Nauka, Moscow, 1985.
- [2] V.N. Gorbuzov and V.Yu. Tushchenco, On R-holomorphic solutions of a system of total differential equations (Russian), *Vestsi Nats. Akad. Navuk Belarusi*, Ser. Fiz.-Mat. Navuk, 1999, No. 3, 124-126.
- [3] V.N. Gorbuzov and V.Yu. Tushchenco, R-holomorphic solutions of a total differential equation, *Differential Equations* 35 (1999), No. 4, 447-452.
- [4] I.V. Gaishun, *Completely solvable Multidimensional differential equations* (Russian), Nauka and Texnika, Minsk, 1983.
- [5] V.N. Gorbuzov, *Integrals of differential systems* (Russian), Grodno State University, Grodno, 2006.
- [6] V.N. Gorbuzov, Autonomy of a system of equations in total differentials, *Differential Equations* 34 (1998), No. 2, 153-160.
- [7] G. Darboux, Mémoire sur les équations différentielles algébriques du premier ordre et du premier degré, *Bull. Sc. Math.*, Vol. 2 (1878), 60-96, 123-144, 151-200.
- [8] V.N. Gorbuzov, *Integrals of systems of equations in total differentials* (Russian), Grodno State University, Grodno, 2005.
- [9] V.N. Gorbuzov, Particular integrals of real autonomous polynomial systems of exact differential equations (Russian), *J. Differential equations and control processes*, 2000, No. 2, 1-36 (<http://www.neva.ru/journal>).

- [10] V.N. Gorbuzov and A.F. Pronevich, Spectral method of jacobian systems integral basic in partial equations construction (Russian), *J. Differential equations and control processes*, 2001, No. 3, 17-45 (<http://www.neva.ru/journal>).
- [11] A.F. Pronevich, Autonomous integrals of linear total differential systems (Russian), Deponent VINITI of *Differentsial Uravneniya* 02.10.2002, No. 1667-B2002, 24 p.
- [12] V.N. Gorbuzov and A.F. Pronevich, Building of the integrals of linear differential systems (Russian), *Vestnik of the Grodno State Univ.*, 2003, Ser. 2, No. 2(22), 50-60.
- [13] A.F. Pronevich, Integrals of linear multidimensional system of non-derogatory matrix structure (Russian), *Mathematical research*, Vol. 10 (2003), 143-152.
- [14] V.N. Gorbuzov and A.F. Pronevich, Integrals of R-linear systems of exact differential (Russian), *Dokl. Akad. Nauk Belarusi* 48 (2004), No. 1, 49-52.
- [15] A.F. Pronevich, Integrals of partial differential systems with R-linear coefficients (Russian), *Vestnik of the Yanka Kupala Grodno State Univ*, 2005, Ser. 2, No. 1(31), 45-52.
- [16] A.F. Pranevich, *R-differentiate integrals of exact differential systems* (Russian), Dissert. Kand. fiz-mat. nauk (Ph.D Thesis), Grodno State Univ., Grodno, 2005.
- [17] A.F. Pranevich, Building of first integrals for linear non-homogeneous system of ordinary differential equations with constant coefficients (Russian), *Vestnik of the Yanka Kupala Grodno State Univ.*, 2008, Ser. 2, No. 2(68), 5-10.
- [18] C. Christopher and J. Llibre, Algebraic aspects of integrability for polynomial systems, *Qualitative Theory of Dynamical Systems*, 1999, No. 1, 71-95.
- [19] M.V. Dolov and S.A. Chistyakova, On the structure of the general solution and integrating factor in a neighborhood of a simple singular point, *Differential Equations*, 37 (2001), No.5, 747-750.
- [20] N.P. Erugin, *A book for reading the general course of differential equations* (Russian), Nauka and Technika, Minsk, 1972.
- [21] M. Falconi and J. Llibre, $n - 1$ independent first integrals for linear differential systems in $\mathbb{R}^n$ and $\mathbb{C}^n$, *Qualitative theory of dynamical systems*, 2004, No. 4, 233-254.
- [22] S.D. Furta, On non-integrability of general systems of differential equations, *Z. Angew. Math. Phys.*, 1996, No. 47, 112-131.
- [23] V.N. Gorbuzov, Autonomous integrals and Jacobi last factors for systems of ordinary differential equations, *Differential Equations* 30 (1994), No. 6, 868-875.
- [24] V.N.Gorbuzov and S.N.Daranchuk, A basis of the autonomous first integrals of the Jacobi-Fourier's system (Russian), *Vestnik of the Belarusian State Univ.*, 2005, No.3, 70-74.
- [25] V.N. Gorbuzov and P.B. Pauliuchyk, On the solutions, integrals and limit cycles of n -Darboux's system (Russian), *J. Differential equations and control processes*, 2002, No. 2, 26-46 (<http://www.neva.ru/journal>).
- [26] V. N. Gorbuzov and V. Yu. Tyshchenko, Particular integrals of systems of ordinary differential equations, *Mat. Sb.* 183 (1992), No. 3, 76-94.
- [27] A. Goriely, Integrability, partial integrability and nonintegrability for systems of ordinary differential equations, *J. Math. Phys.* 37 (1996), 1871-1893.
- [28] A. Goriely, A brief history of Kovalevskaya exponents and modern developments, *Regular and Chaotic Dynamics*, Vol. 5 (2000), No. 1, 3-15.

- [29] B. Grammaticos, J. Moulin Ollagnier, A. Ramani, J.-M. Strelcyn, and S. Wojciechowski, Integrals of quadratic ordinary differential equations in $\mathbb{R}^3$: The Lotka-Volterra system, *Physica A* 163 (1990), 683 -722.
- [30] A.I. Jablonskii, Algebraic integrals of system of differential equations, *Differential Equations* 6 (1970), No. 11, 1326-1333.
- [31] C.G.J. Jacobi, Theoria nova multiplicatoris systemati aequationum differentialium vulgarium applicandi, *J. für reine und angew. Math.* Vol. 27 (1844), 199-268; Vol. 29 (1845), 213-279, 333-376.
- [32] J.P. Jouanolou, *Equations de Pfaff algébriques*, Springer-Verlag, New York, 1979.
- [33] B.M. Koialovich, *Researches on the differential equations $ydy - ydx = Rdx$* (Russian), Sankt Peterburg, 1894.
- [34] A.N. Korkine, Thoughts about multipliers of differential equations of first degree (Russian), *Math. Sbornik* 24 (1904), 194-350 and 351-416.
- [35] S. Kowalevski, Sur le probleme de la rotation d'un corps solide autour d'un point fixe, *Acta Mathematica* 12 (1889), 177-232.
- [36] V.V. Kozlov and D.V. Treschchev, Kovalevskya numbers of generalized Toda chains, *Math. Notes* 46 (1989), 840-848.
- [37] M.N. Lagutinskii, *Partial Algebraic Integrals* (Russian), Kharkov, 1908.
- [38] K.Ja. Latysheva, On the works of V.P. Jermakov on the theory of differential equations (Russian), *Istoriko-Matematicheskije Issledovanija* 9, Gostekhizdat, Moscow (1956), 691-722.
- [39] P.D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Commun. Pure Appl. Math.*, 1968, No. 221, 467-490.
- [40] J. Liouville, Mémoire sur l'intégration d'une classe d'équations différentielles du second ordre en quantités finies explicites, *J. math. pures et appl.*, Vol. 4 (1839), 423-456.
- [41] A.J. Maciejewski, About algebraic integrability and non-integrability of ordinary differential equations, *Chaos, Solitons, Fractals* 9 (1998), 51-65.
- [42] F. Minding, Beiträge zur Integration der Differentialgleichungen erster Ordnung, *Mem. de l'Acad. des Sci. de St.-Petersbourg VII-me série*, Vol. 5 (1862), No. 1, 1-95.
- [43] V.I. Mironenko, *Linear dependence of functions along solution of differential equations* (Russian), Belarusian State University, Minsk, 1981.
- [44] D. Morduhai-Boltovskoi, *On integration of linear differential equations in finite terms* (Russian), Warsaw, 1910.
- [45] J. Moulin Ollagnier and J.-M. Strelcyn, On first integrals of linear systems, Frobenius integrability theorem and linear representations of Lie algebras, in Bifurcations of Planar Vector Fields, Proceedings, Luminy 1989, J.-P. Francoise and R. Roussarie (Eds.), *Lecture Notes in Mathematics* 1455, Springer-Verlag, Berlin, Heidelberg, New-York, 1991.
- [46] A.Nowicki, On the nonexistence of rational first integrals for systems of linear differential equations, *Linear Algebra Appl.* 235 (1996), 107-120.
- [47] M.J. Prellé and M.F. Singer, Elementary first integrals of differential equations, *Trans. Amer. Math. Soc.* 279 (1983), 215-229.

- [48] M.F. Singer, Liouvillian first integrals of differential equations, *Trans. Amer. Math. Soc.* 333 (1992), 673-688.
- [49] D. Schlomiuk, Algebraic particular integrals, integrability and the problem of the center, *Trans. Amer. Math. Soc.*, 338 (1993), No. 2, 799-841.
- [50] D. Schlomiuk and N. Vulpe, Geometry of quadratic differential systems in the neighbourhood of the line at infinity, *Preprint of the Cornell University*, arXiv:math/0405026v1 [math.DS] 3 May 2004.
- [51] A. Tsygvintsev, On the existence of polynomial first integrals of quadratic homogeneous systems of ordinary differential equations, *J. Phys. A: Math. Gen.* 34 (2001), 2185-2193.
- [52] J. A. Weil, *Constantes et polynomes de Darboux en algebre differentielle: applications aux systemes differentiels lineaires* (Ph.D Thesis), Ecole polytechnique, 1995.
- [53] H. Yoshida, Necessary conditions for the existence of algebraic first integrals I, II, *Celest. Mech.* 31 (1983), 363-379 and 381-399.
- [54] S.L. Ziglin, Branching of solutions and nonexistence of first integrals in Hamiltonian mechanics, *Functional Anal. Appl.* 16 (1983), 181-189.
- [55] D.V. Buslyuk, Integrals and last multipliers of partial differential systems, *Differential Equations* 35 (1999), No. 3, 421-422.
- [56] N.M. Gjunter, *Integrirovanie uravnenij v chastnyh proizvodnyh 1-go porjadka* (Russian), GTTI, Moscow-Leningrad, 1934.
- [57] V.N. Gorbuzov and S.N. Daranchuk, The integrals and last multipliers of one class of partial differential systems (Russian), *J. Differential equations and control processes*, 2007, No. 4, 1-16 (<http://www.neva.ru/journal>).
- [58] E. Kamke, *Differentialgleichungen: Lösungsmethoden und Lösungen*, II, Partielle Differentialgleichungen Erster Ordnung für eine gesuchte Funktion, Akad. Verlagsgesellschaft Geest & Portig, Leipzig, 1965.
- [59] A.D. Polanin, V.F. Zaitsev, and A. Moussiaux, *Handbook of first order partial differential equations*, Taylor & Francis, London, 2002.
- [60] V.V. Amel'kin, *Autonomous and linear many-dimensional differential equations* (Russian), Universitetskoe, Minsk, 1985.
- [61] A.R. Forsyth, *Theory of differential equations*, Vol. 1-6, Cambridge University Press (1890-1906), reprinted by Dover Public., New York, 1959.
- [62] V.N. Gorbuzov and A.F. Pranevich, Autonomy and cylindricity of R-differentiable integrals for systems in total differentials (Russian), *J. Differential equations and control processes*, 2008, No. 1, 35-49 (<http://www.neva.ru/journal>).
- [63] E. Goursat, *Cours d'analyse mathematique*, Vol. 2, Paris (4th. ed. 1924). English translation: *A Course of Mathematical Analysis*, Vol II, Part Two: Differential Equations, Dover Public., New York, 1959.
- [64] F.R. Gantmacher, *The theory of matrices* (Russian), Nauka, Moscow, 1988.
- [65] R.A. Horn and C.R. Johnson, *Matrix analysis*, Cambridge Univ. Press, Cambridge, 1990.